

The author(s) shown below used Federal funds provided by the U.S. Department of Justice and prepared the following final report:

**Document Title: Prevalence and Consequences of Child
Victimization: Results from the National Survey
of Adolescents, Final Report**

**Author(s): Dean G. Kilpatrick Ph.D. ; Benjamin E. Saunders
Ph.D.**

Document No.: 181028

Date Received: 03/29/2000

Award Number: 93-IJ-CX-0023

This report has not been published by the U.S. Department of Justice. To provide better customer service, NCJRS has made this Federally-funded grant final report available electronically in addition to traditional paper copies.

**Opinions or points of view expressed are those
of the author(s) and do not necessarily reflect
the official position or policies of the U.S.
Department of Justice.**

***Prevalence and Consequences of Child Victimization:
Results from the
National Survey of Adolescents***

Final Report

Dean G. Kilpatrick, Ph.D.
Benjamin E. Saunders, Ph.D.
Principal Investigators

Medical University of South Carolina
National Crime Victims Research and Treatment Center
Department of Psychiatry and Behavioral Sciences
171 Ashley Avenue
Charleston, South Carolina 29425-0742
(803) 792-2945



November, 1997

This project was funded by the U.S. Department of Justice, Office of Justice Programs, National Institute of Justice, Grant No. 93-IJ-CX-0023. Opinions expressed in this report do not necessarily represent the official position, policies, or procedures of the U.S. Department of Justice.

Acknowledgments

This project would not have been possible without the contributions of a number of people, each of whom we would like to thank. Several members of the National Crime Victims Research and Treatment Center (NCVRTC) faculty made major contributions to the project design, data analyses, and interpretation of findings. These include Drs. Heidi Resnick, Connie Best, Julie Crouch, and Ron Acierno. We are extremely grateful for their assistance. A number of NCVRTC postdoctoral fellows also contributed to the project including Drs. Elizabeth Letourneau and Daniel Smith. Members of the Advisory Committee provided sage advice and excellent guidance. We are grateful to Lucy Berliner and to Drs. Del Elliott, David Finklehor, Scott Henggler, and Cathy Widom for agreeing to serve as Advisory Committee members and for the outstanding job they did. The project would not have been possible without the extraordinary contributions of Dr. John Boyle and his colleagues at Schulman, Ronca, & Bucavalas, Inc. John's efforts and SRBI's professionalism were primarily responsible for the NSA's excellent response and completion rate as well as for the quality of the interview data that were gathered. We also are grateful for all of the help and encouragement we received from our National Institute of Justice grant monitor, Bernie Auchter. Finally, we would like to thank NCVRTC staff members Vickey Cornelison and Dori Long for all their assistance in preparing this document.

Table of Contents

I.	INTRODUCTION	1
A.	Specific Aims	1
II.	STATEMENT OF HYPOTHESES	3
III.	REVIEW OF LITERATURE	4
IV.	RESEARCH DESIGN AND METHODS	9
A.	Research Participants	9
B.	Sample Development Strategy	9
C.	Recruitment and Interview Procedures	10
1.	<u>Initial interview with the parent or guardian</u>	10
2.	<u>Selection of the Adolescent Respondent within Households</u>	10
3.	<u>Interviewing the Designated Adolescent</u>	11
4.	<u>Verification of Survey Authenticity</u>	11
5.	<u>Incentives</u>	12
6.	<u>Training of Interviewers</u>	12
7.	<u>Rationale for Conducting the Study by Telephone</u>	12
D.	Survey Instruments and Measurement of Key Constructs	13
1.	<u>Biographic/Demographic Characteristics</u>	13
2.	<u>Family History of Substance Abuse</u>	13
3.	<u>Victimization History</u>	14
4.	<u>Posttraumatic Stress Disorder (PTSD)</u>	14
5.	<u>Substance Use</u>	15
6.	<u>Substance Dependence and Abuse</u>	15
a.	<i>Lifetime Substance Dependence</i>	15
c.	<i>Current Substance Dependence</i>	16
d.	<i>Lifetime Substance Abuse</i>	16
e.	<i>Current Substance Abuse</i>	16
7.	<u>Delinquent Behaviors</u>	16
E.	Sample Disposition and Participation Rates:	16
F.	Demographic Characteristics of Parent and Adolescent Samples	17
1.	<u>Parent Sample</u>	17
2.	<u>The Adolescent Sample</u>	17
V.	RESULTS	19
A.	Overview of Data Analytic Strategy	19
B.	Descriptive Results	19
1.	<u>Family History Variables</u>	19
2.	<u>Personal Victimization</u>	19

2.	<u>Sexual Assault</u>	19
	(1) <i>Lifetime Prevalence</i>	19
	(2) <i>Age at Time of the Assault</i>	20
	(3) <i>Relationship to the Perpetrator</i>	20
	(4) <i>Location of Assault</i>	20
	(5) <i>Life Threat and Physical Injury</i>	20
	(6) <i>Reporting of Cases to Authorities</i>	21
	b. <i>Physical Assault</i>	21
	(1) <i>Lifetime Prevalence</i>	21
	(2) <i>Age at Time of Assault</i>	21
	(3) <i>Relationship to the Perpetrator</i>	21
	(4) <i>Location of Assault</i>	21
	(5) <i>Life Threat and Physical Injury</i>	21
	(6) <i>Reporting to Police or Other Authorities</i>	22
	c. <i>Physically Abusive Punishment</i>	22
3.	<u>Witnessing Violence</u>	22
4.	<u>Posttraumatic Stress Disorder</u>	22
5.	<u>Substance Use/Abuse Dependence</u>	22
	a. <i>Past-Year and Lifetime Alcohol, Marijuana, and Hard Drug Use</i>	22
	b. <i>Past-Year and Lifetime Alcohol, Marijuana, and Hard Drug Abuse or Dependence</i>	23
	c. <i>Past-Year and Lifetime Abuse or Dependence of Any Substance</i>	23
6.	<u>Delinquent Behavior</u>	23
C.	<u>Rates of Personal Victimization, Witnessing Violence, PTSD, Substance Use/Abuse/Dependence, and Delinquency Variables by Demographic Characteristics</u>	23
1.	<u>Introduction</u>	23
2.	<u>Sexual Assault</u>	23
	a. <i>Gender</i>	23
	b. <i>Age</i>	24
	c. <i>Race/Ethnicity</i>	24
3.	<u>Physical Assault</u>	24
	a. <i>Gender</i>	24
	b. <i>Age</i>	24
	c. <i>Race/Ethnicity</i>	24
4.	<u>Physically Abusive Punishment</u>	24
	a. <i>Gender</i>	24
	b. <i>Age</i>	24
	c. <i>Race/Ethnicity</i>	25
5.	<u>Witnessing Violence</u>	25
	a. <i>Gender</i>	25
	b. <i>Age</i>	25
	c. <i>Race/Ethnicity</i>	25
6.	<u>PTSD</u>	25

	a.	<i>Gender</i>	25
	b.	<i>Age</i>	25
	c.	<i>Race/Ethnicity</i>	25
7.		<u>Substance Use</u>	26
	a.	<i>Gender</i>	26
	b.	<i>Age</i>	26
	c.	<i>Race/Ethnicity</i>	26
8.		<u>Current Substance Abuse/Dependence</u>	26
	a.	<i>Gender</i>	27
	b.	<i>Age</i>	27
	c.	<i>Race/Ethnicity</i>	27
9.		<u>Delinquent Behaviors</u>	27
	a.	<i>Gender</i>	27
	b.	<i>Age</i>	27
	c.	<i>Race/Ethnicity</i>	27
D.		Univariate Relationships Between Victimization Variables and Dependent Variables (PTSD, Any Substance Abuse/Dependence, Past-Year Delinquency)	27
	1.	<u>Posttraumatic Stress Disorder (Current and Lifetime)</u>	27
		a. <i>Sexual Assault and PTSD</i>	27
		b. <i>Physical Assault or Physically Abusive Punishment and PTSD</i>	28
		c. <i>Witnessed Violence and PTSD</i>	28
	2.	<u>Any Substance Abuse/Dependence (Current and Lifetime)</u>	28
		a. <i>Sexual Assault and any Substance Abuse/Dependence</i>	28
		b. <i>Physical Assault and any Substance Abuse/Dependence</i>	29
		c. <i>Physically Abusive Punishment and any Substance Abuse/Dependence</i>	29
		d. <i>Witnessed Violence and any Substance Abuse/Dependence</i>	29
	3.	<u>Past-Year Delinquency</u>	30
		a. <i>Sexual Assault and Past Year Delinquency</i>	30
		b. <i>Physical Assault and Past Year Delinquency</i>	30
		c. <i>Physically Abusive Punishment and Past Year Delinquency</i>	30
		d. <i>Witnessed Violence and Past Year Delinquency</i>	30
E.		Multivariate Analyses of the Relationship Between Violence Exposure and Posttraumatic Stress Disorder	30
	1.	<u>Overview of Analyses</u>	30
	2.	<u>Results of Univariate Analyses</u>	31
	3.	<u>Results of Multivariate Logistic Regression Analysis</u>	31
F.		Multivariate Analyses of the Relationship Between Violence Exposure and Past Year Substance Abuse/Dependence	33
	1.	<u>Overview of Analyses</u>	33
	2.	<u>Results of Univariate Analyses</u>	34
		a. <i>Alcohol Abuse/Dependence</i>	34
		b. <i>Marijuana Abuse Dependence</i>	34
		c. <i>Hard Drug Abuse/Dependence</i>	34
	3.	<u>Multivariate Logistic Regression Analyses</u>	34

a.	<i>Alcohol Abuse/Dependence</i>	34
b.	<i>Marijuana Abuse/Dependence</i>	35
c.	<i>Hard Drug Abuse/Dependence</i>	36
4.	<u>Timing of Exposure to Violence and First Use of Alcohol or Drug</u>	36
G.	Multivariate Analyses of the Relationship Between Violence Exposure and Past Year Delinquency	37
1.	<u>Multivariate analysis of delinquency correlates</u>	37
H.	Prevalence Summary and Population Estimates of Critical Study Variables ...	41
VI.	DISCUSSION	43
VII.	RECOMMENDATIONS FOR RESEARCH, POLICY, AND PRACTICE	47
A.	Recommendations for Research	47
B.	Recommendations for Policy	48
C.	Recommendations for Practice	49
VIII.	REFERENCES	51
Table 1:	<u>Demographic Characteristics of Parent/Guardian Sample</u>	57
Table 2:	<u>Demographic Characteristics of Adolescent Sample</u>	60
Table 3:	<u>Lifetime Prevalence of Types of Sexual Contact by Gender</u>	62
Table 4:	<u>Lifetime Prevalence of Physical Assault by Gender</u>	63
Table 5:	<u>Lifetime Prevalence of Physically Abusive Punishment by Gender</u>	64
Table 6:	<u>Lifetime Prevalence of Witnessing Violence by Gender</u>	65
Table 7:	<u>Initial and Final Model Odds of Lifetime PTSD as a Function of Demographic Variables, Family Substance Use, and Sexual Assault, Physical Assault, and Witnessed Violence: Hierarchical Logistic Regression Analyses</u>	66
Table 8:	<u>Univariate Odds of Past-Year Alcohol Abuse/Dependence, Marijuana Abuse/Dependence, and Hard Drug Abuse/Dependence as a Function of Demographic Variables, Familial Problem Alcohol Use, Familial Problem Drug Use, Physical Abuse/Assault, Sexual Assault, Witnessed Violence, and Current PTSD: Chi Square Analyses</u>	67
Table 9:	<u>Initial and Final Model Odds of Past-Year Alcohol Abuse/Dependence, Marijuana Abuse/Dependence, and Hard Drug Abuse/Dependence as a Function of Demographic Variables, Familial Problem Alcohol Use, Familial Problem Drug Use, Physical Abuse/Assault, Sexual Assault, Witnessed Violence, and Current PTSD: Hierarchical Logistic Regression Analyses</u>	69

Table 10:	<u>Hierarchical Logistic Regression: Initial and Final Odds Ratio for Prediction of Past-year Delinquency for Male and Female Adolescents</u>	71
Table 11:	<u>Predictive Models of Past-year Index Offenses for White and African-American Males and Females</u>	72
Table 12:	<u>Proportion of Male Youths Who Committed an Index Offense in the Past Year with Risk Factor Combination</u>	73
Table 13:	<u>Proportion of Female Youths Who Committed an Index Offense in the Past Year with Risk Factor Combination</u>	74
Table 14:	<u>Population Estimates of Victimization and PTSD</u>	75
Figure 1:	Age at Time of Sexual Assault	76
Figure 2:	Relationship Between Victim and Perpetrator in Sexual Assault Cases	77
Figure 3:	Location of Sexual Assaults	78
Figure 4:	Life Threat and Degree of Physical Injury Sustained During Sexual Assault	79
Figure 5:	Reporting of Sexual Assault to Authorities	80
Figure 6:	Victims Age at Time of Physical Assault	81
Figure 7:	Relationship Between Physical Assault Victims and their Perpetrators	82
Figure 8:	Location of Physical Assault	83
Figure 9:	Life Threat and Degree of Physical Injury Sustained During Physical Assault	84
Figure 10:	Reporting of Physical Assault to Authorities	85
Figure 11:	Lifetime Prevalence of Witnessing Violence	86
Figure 12:	Lifetime Prevalence of Sexual Assault by Age Cohort	87
Figure 13:	Lifetime Prevalence of Physical Assault by Age Cohort	88
Figure 14:	Victims Age at Time of Physically Abusive Punishment by Age Cohort	89
Figure 15:	Age at Time Witnessed Violence by Age Cohort	90
Figure 16:	Prevalence Rates of Lifetime and Current PTSD by Age Cohort	91
Figure 17:	Lifetime and Past Year Heavy Alcohol Use by Age Cohort	92
Figure 18:	Lifetime and Past Year Use of Illicit Drugs by Age Cohort	93
Figure 19:	Past Year Rates of Alcohol, Marijuana, and Hard Drug Abuse/Dependence	94
Figure 20:	Lifetime and Past Year Delinquent Offense by Age Cohorts	95
Figure 21:	Percentage of Male and Female Adolescents with Lifetime PTSD by Number of Sexual Assaults Experienced	96
Figure 22:	Percentage of Male and Female Adolescents with Lifetime PTSD by Number of Physical Assaults Experienced	97
Figure 23:	Percentage of Male and Female Adolescents with Lifetime PTSD by Number of Incidents of Violence Witnessed	98
Figure 24:	Rates of Past Year Any Substance Abuse/Dependence by Victimization Risk Factors and Gender	99
Figure 25:	Past Year Delinquency by Victimization Risk Factors and Gender	100

Figure 26.	Percentage of Male and Female Adolescents with PTSD by Family Members and Alcohol and Drug Problems	101
Figure 27.	Number of Violent Incidents Experienced or Witnessed by Male and Female Adolescents	102
Figure 28:	Past Year Alcohol Abuse/Dependence	103
Figure 29:	Past Year Marijuana Abuse/Dependence	104
Figure 30:	Past Year Hard Drug Abuse/Dependence	105

I. INTRODUCTION

A. Specific Aims

Violence is a substantial problem for American youth. Past research and recent governmental reports indicate that many children and adolescents have been victims of violence and that a growing number of youth are perpetrators of serious violence (Centers for Disease Control, 1990; National Center on Child Abuse and Neglect, 1994; Finkelhor & Dzuiba-Leatherman, 1994; Bastian & Taylor, 1991; Elliott, 1992). Family members are often the perpetrators of violence against children and adolescents, as well as the targets of their violence. Victimization in childhood has been found to be strongly related to subsequent substance use/abuse/dependence and delinquent behavior (Huizinga, Loeber & Thornberry, 1995; Widom, 1992; Dembo, Williams, Schmeidler, Berry, Wothke, Getreu, Wish, & Christensen, 1992; Kilpatrick, Resnick, Saunders, Best, & Epstein, 1994), as well as a variety of other mental, emotional, and behavioral problems including Post-traumatic Stress Disorder and depression (Browne & Finkelhor, 1986; Beitchman, Zucker, Hood, DaCosta, Akman, & Cassavia, 1992; Saunders, Villepontoux, Lipovsky, Kilpatrick & Veronen, 1992; Murphy, Amick-MuMullen, Kilpatrick, Haskett, Veronen, Best, & Saunders, 1988; Lipovsky, Saunders & Murphy, 1989). In general, studies have examined "family violence" and ignored other victimization experiences, or assessed "criminal violence" and disregarded violence perpetrated by family members. This arbitrary distinction has given rise to two separate scientific literatures, created a false dichotomy between family and nonfamily victimization and violence experiences, and discouraged a more complete and integrated view of assessing the etiology and consequences of victimization and violence. In response to this problem, this study will assess both familial and nonfamilial violence.

A substantial number of studies have examined different aspects of the relationships between childhood victimization experiences, trauma-related mental health problems, substance use/abuse/dependency, and delinquent behavior. However, many of these studies suffer from critical conceptual and methodological problems that limit their utility. More important, most studies to date have examined only selected relationships between these constructs. Some have tested the relationships between victimization and mental health effects; others the relationships between substance use/abuse/dependency and delinquency; and still others the relationships between childhood victimization and delinquency. However, none have tested a comprehensive conceptual framework linking all of these complex relationships into an explanatory model. Most studies also suffer from one or more serious methodological problems such as using nonrepresentative samples, not thoroughly assessing participants for a history of victimization and trauma, ignoring or poorly measuring mental health problems, or not adequately examining potential gender and racial/ethnic differences. In addition, most studies use retrospective rather than longitudinal designs or rely primarily on official case records rather than directly assessing individuals.

The National Survey of Adolescents, funded by the National Institute of Justice (grant number 93-IJ-CX-0023) addressed several of these conceptual and methodological problems. The goal of this study was to test specific hypotheses generated by a theoretically and

empirically constructed conceptual framework illustrating the relationships between serious victimization experiences, the mental health effects of victimization, substance use/abuse and delinquency behavior in addition to demographic and important background variables. adolescents were assessed for a history of sexual assault, physical assault, harsh physical discipline, witnessing violent events, Posttraumatic Stress Disorder (PTSD), depression, substance use/abuse/dependence, and commission of index delinquency offenses. The NSA was a telephone survey of a nationally representative sample of 4,023 American youth between the ages of 12 and 17 living in U. S. households with telephones. Data collection occurred between January and June, 1995.

Specific aims of the NSA were to

1. provide descriptive information about cases of familial and nonfamilial violent assault, delinquent behavior, mental health problems, and substance use, abuse and dependence broken down by age, gender, family income, and racial/ethnic group among American adolescents;
2. test a risk factor model that hypothesizes relationships between violent familial and nonfamilial victimization in childhood and adolescence and risk of PTSD, delinquent behavior, and substance use/abuse/dependence among American adolescents; and
3. examine potential differences between gender and ethnic minority groups in the etiology and consequences of substance use/abuse/dependence and delinquent behavior among American adolescents

II. STATEMENT OF HYPOTHESES

The primary hypothesis of this study is that victimization during childhood and or adolescence increases the risk of developing significant psychological distress and substance use, which in turn increases risk for substance abuse or dependence, delinquent behavior, and subsequent victimization. Specific hypothesis to be tested are:

- H₁: Victimized adolescents (whether by family members or others) will be more likely than nonvictimized adolescents to have high levels of psychological distress.
- H₂: Victimized adolescents (whether by family members or others) will be more likely than nonvictimized adolescents to use alcohol and illicit drugs.
- H₃: Victimized adolescents (whether by family members or others) with high levels of psychological distress will be more likely than victimized adolescents with low levels of psychological distress to have substance use, abuse, and dependency problems.
- H₄: Victimized adolescents (whether by family members or others) will be more likely than nonvictimized adolescents to engage in delinquent behavior.
- H₅: Victimization (whether by family members or others) temporally precedes initial problems with psychological distress, substance abuse, and delinquent behavior.
- H₆: Lifetime and past year prevalence of sexual assault will be higher among girls than among boys; prevalence of physical assault will be higher among boys than among girls; prevalence of witnessing violence will not differ as a function of gender.
- H₇: Prevalence rates of violent victimization (whether by family members or others) will not differ across ethnic/racial groups after controlling for the effects of age, gender, family income, and residential location.
- H₈: Causal pathways for delinquent behavior, substance use, substance dependence, and substance abuse will differ as a function of gender, but will not differ by ethnic/racial status.

III. REVIEW OF LITERATURE

Considerable evidence exists suggesting that both family and nonfamily violence is a major problem for children and adolescents, both in its prevalence and consequences (National Center on Child Abuse and Neglect, 1994; Finkelhor & Dziuba-Leatherman, 1994; Kilpatrick, Edmunds, & Seymour, 1992; Bastian & Taylor, 1991; Finkelhor, Hotaling, Lewis, & Smith, 1990; Gelles & Straus, 1987; McCurdy & Daro, 1993; Saunders et al., 1992; Burnam et al., 1988; Whitaker & Bastian, 1991; Elliott, 1992). A history of violent assault during childhood or adolescence increases risk for a host of major mental health problems such as Post-traumatic Stress Disorder and depression (Ageton, 1983; Browne & Finkelhor, 1986; Beitchman, et al., 1992; Burnam, Stein, Golding, Siegel, Sorenson, Forsythe, & Tefles, 1988; Saunders, et al., 1992; Jaffe, Wolfe & Wilson, 1990; Lanktree, Briere, & Zaidi, 1991), and substance use/abuse/dependency problems (Ageton, 1983; Burnam et al. 1988; Gelles & Straus, 1990; Saunders, Kilpatrick, Lipovsky, Resnick, Best, & Sturgis, 1991; Stein, Golding, Siegel, Burnam, & Sorenson, 1988). Still other evidence suggests that youth victimization history increases risk of involvement with delinquent peers and of subsequent delinquent behavior (Ageton, 1983; Dembo et al. 1992; Straus, 1984; Widom, 1989; Widom, 1992; Huizinga, et al. 1995). Some research shows that involvement with delinquent or deviant peers increases risk of victimization (eg., Ageton, 1983), and that substance use also increases risk of victimization (eg. Kilpatrick et al., in press; Kilpatrick, Resnick, Saunders, Best & Epstein, 1994; Cottler, Compton, Mager, Spitznagel, & Janca, 1992). Other research indicates that there is substantial comorbidity between PTSD and substance use, dependence, and abuse (Cottler et al., 1992; Kessler, Sonnega, Bromet, Hughes & Nelson, 1995). Therefore, the constructs of familial and nonfamilial victimization in childhood or adolescence, trauma-related mental health problems, substance use/abuse/dependency, and delinquent behavior are inter-related in a variety of ways.

Two lines of research with adults confirm that victimization is a risk factor for substance use/dependency/abuse and visa versa. First, epidemiological studies show that substance use disorders are more prevalent in individuals who have a history of criminal victimization (Burnam et al. 1988; Cottler et al. 1992; George & Winfield-Laird, 1986; Helzer, Robins, & McEvoy, 1987; Kilpatrick, 1990; Kulka, Schlenger, Fairbank, Hough, Jordan, Marinar, & Weiss, 1990; Sorenson, Stein, Siegel, Golding, & Burnam, 1987). For example, in a study of 3,125 Los Angeles residents, as part of the NIMH Epidemiologic Catchment Area project, rates of substance abuse or dependence (both alcohol and other drug) were significantly higher among sexual assault victims compared to non-victims (Sorenson et al.).

Second, studies of women seeking treatment for substance use/abuse/dependency problems have high rates of victimization (Brady, Killen, Saladin, Dansky & Becker, 1994; Ladwig & Anderson, 1989; Miller, Downs, Gondoli & Keil, 1987; Miller, Wiczorek & Downs, 1994; Kilpatrick & Resnick, 1994). Both these lines of research confirm that there is a relationship between victimization and substance use/abuse/dependency, but because of the cross-sectional, retrospective nature of most extant studies, it is impossible to establish the temporal or causal sequence of events.

Using student samples and a national sample, Straus (1984) found that parental violence towards teenage children was related to children's rate of crime and violence outside the family in addition, boys raised in families characterized by interparental violence had higher rates of delinquency and aggression towards others outside the family. Finally, witnessing violence itself appears to be associated with later violent behavior (e.g., Lewis, Shanok, Pincus, & Glaser, 1979).

The most methodologically sound study examining the relationships among adolescent victimization, delinquent behavior, substance use/abuse problems, and mental health problems, is the National Youth Survey (NYS) conducted by Delbert Elliott and his associates (Ageton, 1983; Elliott, Huizinga & Ageton, 1985; Elliott, Huizinga, & Menard, 1989). The NYS is a theory-driven longitudinal study of a national probability household sample of 1,725 adolescents between the ages of 11 and 17 at the time of the projects' onset in 1976. Most published work focuses on the first six waves of data collection. The NYS obtained annual self-report estimates of the respondent's frequency of delinquent behavior, drug and alcohol use, substance abuse-related problems, and some mental health problems (primarily major depression). Limited information was obtained about sexual assault from female respondents but not about physical and sexual assault occurring a year or more prior to the first wave of the study.

Two issues concerning the NYS findings are particularly relevant to this project. First, Ageton (1983) concluded that "engaging in delinquent behavior and being a part of a delinquent network influence the risk of being sexually assaulted". However, the study design (which did not measure sexual assaults occurring in childhood, i.e., prior to age 11) and the relatively weak sexual assault screening questions make it impossible to determine whether a history of sexual assault in childhood might have preceded the delinquent behavior and/or exposure to delinquent peers. Second, the NYS found a high degree of overlap between delinquent behavior, exposure to delinquent peers, and substance use/abuse problems (Elliott et al., 1985). In fact, common etiological pathways were found for illegal drug use and delinquency, with prior drug use and/or delinquency as well as exposure to delinquent peers being the best predictors of current drug use and/or delinquency. Results obtained with longer follow-up and adding mental health problems to the assessment were more complex (Elliott, Huizinga, & Menard, 1988). As adolescents become young adults, mental health problems tend to increase, as do drug abuse problems, but delinquency problems tend to decrease. However, the subset of adolescents with all three problems were found to be the most likely to be arrested (Elliott et al., 1988). Given Widom's (1992) findings about child victimization increasing risk of delinquency and adult criminal behavior, the lack of good data on child victimization in the NYS is a real limitation in an otherwise exemplary study. Likewise, the NYS did not measure PTSD, a mental health problem that occurs frequently after victimization, and that has been shown to increase risk for alcohol and drug use problems (Kilpatrick, 1990; Kilpatrick et al., 1992).

In summary, the extant literature provides substantial support for the notion that child victimization is a risk factor for delinquency, substance use problems, and mental health problems such as PTSD and depression. However, previous studies left gaps in knowledge because they did not: 1) use a large national probability sample of adolescents of both genders with a substantial representation of ethnic minority groups; 2) measure baseline history of a

broad range of familial and nonfamilial childhood victimization experiences; 3) assess the spectrum of substance use ranging from tobacco to hard drugs; 4) assess important potentially comorbid mental disorders such as PTSD and depression; 5) examine potential consequences of victimization and substance use such as substance abuse/dependency, delinquent behavior, and risk of revictimization; 6) test alternative pathways for development of substance use; and 8) base hypothesis and design on theory. The present study was specifically designed to fill this gap.

IV RESEARCH DESIGN AND METHODS

A Research Participants:

The NSA sample consisted of two subsamples, a national probability household sample of 3,161 adolescents and a probability oversample of 862 adolescents residing in central city areas of the United States. Eligible for selection were all adolescents between the ages of 12 and 17 living in households with telephones in the United States, who resided with a parent or guardian, and who could converse in English or Spanish. The only adolescents potentially excluded from the study were those residing in institutional settings, in households without a parent or guardian (e.g., emancipated minors, married adolescents living on their own) or in households without telephones; those who did not speak English or Spanish, and those whose parents did not give permission for their adolescent to be interviewed. According to the 1990 census, only 5% of all U.S. households did not have telephones at any one point in time. Based on the results of a large RDD survey of adolescents sampled using similar methodology conducted by the applicants (Boyle & Kilpatrick, 1993), less than 2% of otherwise eligible adolescents do not speak English or Spanish. Therefore, we estimate that the sampling frame provided coverage for about 93% of U. S. adolescents living in households with parents or guardians and should be highly representative of U. S. adolescents living in households with parents or guardians.

In addition to the adolescent participants in the NSA, one parent or guardian in each household was interviewed briefly as will be described subsequently. The primary purpose of these interviews was to establish rapport and to obtain permission to interview the targeted adolescent. Because the parent or guardian interviews were conducted prior to the adolescent interviews, the 4,023 participants in the parent sample were also selected from a national probability sample of households and a probability oversample of central city households. Like the case with adolescents, parents were eligible if they spoke English or Spanish.

All sample selection and interviewing was done by Schulman, Ronca, and Bucuvalas Inc. (SRBI), a New York-based survey research firm.

B. Sample Development Strategy:

To construct the initial national probability sample, the NSA used a multi-stage, stratified, area probability, random digit dialing (RDD) sampling procedure that had four steps. First, the U.S. was stratified geographically by census region and a population-based subsample allocation was developed for each geographic stratum. In other words, the number of households drawn for the sample from each geographic stratum was allocated in proportion to the actual distribution of the population residing within each stratum according to the most recent census estimates. Geographical stratification was used because the precision of sample estimates generally are improved by stratification. Hence, the population of the United States was stratified by census region. Specifically, the regional stratification of the sample was divided into the nine census regions as follows:

<i>New England:</i>	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut
<i>Middle Atlantic:</i>	New York, New Jersey, and Pennsylvania
<i>East North Central:</i>	Ohio, Indiana, Illinois, Michigan, and Wisconsin
<i>West North Central:</i>	Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas
<i>South Atlantic:</i>	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, and Florida
<i>East South Central:</i>	Kentucky, Tennessee, Alabama, and Mississippi
<i>West South Central:</i>	Arkansas, Louisiana, Oklahoma, and Texas
<i>Mountain</i>	Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, and Nevada
<i>Pacific:</i>	Washington, Oregon, California, Alaska, and Hawaii

The estimated distribution of the adolescent population by stratum was calculated on the basis of the Projections of the Population of States by Age, Sex and Race: 1988 to 2010 (Current Population Reports, P-25, No. 1017, 1988).

In the second step, telephone banks within each geographic stratum were systematically selected utilizing the comprehensive database of working telephone banks maintained by SRBI. Third, random digit dialing (RDD) was used to sample telephone households within the telephone banks selected in the second stage. RDD was used to locate currently working, residential household telephone numbers with eligible respondents. Non-working numbers and non-household (e.g., business) numbers were immediately replaced by other RDD numbers selected within the same stratum in the same fashion as the initial number. Non-answering numbers were recalled four times before being replaced. In the fourth step, an adult respondent in each household was screened to determine if there were any adolescents, 12 to 17, currently living in the household or if any other adolescent had lived in the household at least four months within the previous year.

In households with multiple eligible adolescents, a systematic selection was made to determine which eligible individual would be designated as the respondent. These procedures yielded a relatively unbiased sample of 3,161 adolescents from which valid generalizations can be made to the total population, within specified limits of expected sampling variability.

Construction of the central city oversample followed these same procedures except for the initial geographical stratification step. This step was replaced by using the Census classification of counties by types of place (i.e., central city) and specifying our target population as households located within these urban counties. These were then systematically sampled. The SRBI database of working telephone exchanges and banks of telephone numbers includes county designation. These then were systematically sampled within the selected urban counties. The RDD step was limited to these selected exchanges and banks within the selected urban counties. The third and fourth stages of the sampling procedure (for eligible households and adolescents) for the central city oversample were the same used in the national probability sample.

C. **Recruitment and Interview Procedures:**

1. **Initial interview with the parent or guardian:**

After determining that the household contained one or more eligible adolescents, interviewers asked to speak to a parent or a guardian. Interviews with adolescents were attempted only after a parent or guardian was interviewed and agreed to permit the designated adolescent to be interviewed. If the person with whom the household screening was conducted was not the parent or guardian, then the parent/guardian portion of the interview began with a recap of the general introduction. Then, the parent or guardian was provided with additional relevant information about the study, including:

- How their household was been selected (as part of a national random sample of households with children);
- The length of the parent/guardian portion of the interview (about 10 minutes)
- The sponsorship of the study;
- The general purpose of the study;
- The voluntary nature of the study, and the confidentiality of their responses.

Parent/guardians were provided the opportunity to call a toll-free number to confirm the authenticity of the study. Finally, parents/guardians were asked to grant permission for their adolescent to be interviewed.

The primary objectives of the parent/guardian interview were to inform the parent about the study, secure permission to interview the designated adolescent, and to ensure the collection of comparative data to examine potential non-response bias from households without adolescent participation. In addition to demographics, adult respondents were asked about their concerns with the safety of and potential violence toward their children; their child's past experience with violence; and their (parent's) experience with violence.

All interviews with both parents and adolescents were conducted using Computer-Assisted Telephone Interviewing (CATI) technology. All interviews were transcribed into the SRBI CATI system, and all interviewers used the CATI system. This technology has several advantages. It is better able to handle complex skip patterns and question ordering in complicated interview schedules such as those used in the NSA. It insures that all questions will be asked since interviewers cannot proceed without making responses. CATI interviews typically take less time and result in less respondent fatigue, increasing compliance and reducing termination rates. It also greatly reduces the time for data reduction and cleaning.

2. **Selection of the Adolescent Respondent within Households:**

The sample construction described earlier yields a population-based sample of households with adolescent children. During the parent guardian section of the interview, the parent was asked to enumerate the ages and sex of all eligible children in the household. In the case of households with multiple adolescent children, the adolescent with the most recent

birthday was selected for interview. The "most recent birthday" technique is a common and accepted one for randomly selecting respondents within eligible age cohorts of household members. This method has been demonstrated to be technically equivalent or superior to other respondent selection techniques, and involves less respondent burden.

3. Interviewing the Designated Adolescent:

Whenever possible, adolescents were interviewed immediately following the parent/guardian interviews. Otherwise, appointments were scheduled when possible or blind callbacks at different times of day and days of the week were made. Unlimited callbacks were made throughout the field period before a case was abandoned. The introduction of the study to the designated adolescent included the following:

- What the survey is about (dangerous things that happen in their school, neighborhood, and family);
- Why the survey is being conducted;
- How they had been selected (by dialing random phone numbers);
- The types of questions they would be asked (dangerous situations they may have been faced with and personal situations where they may have been threatened);
- Assurance of confidentiality (nothing they said would be told to their parents, school, or anyone else);
- How long the interview would take (about half an hour);
- They don't have to answer any question that they don't want to answer;
- They can terminate the interview at anytime by simply hanging up.

Assent to proceed with the interview was obtained from each adolescent after the explanation.

An important concern was whether the adolescent could answer interview questions freely and in private. Two steps were taken to increase the likelihood that adolescents could answer questions in an open and honest manner with a reasonable degree of privacy. First, the interviewer specifically asked if the adolescent was in a situation where they could be assured of privacy and could answer in an open manner. If the adolescent indicated they could not, the interviewer offered to call back at another time when privacy was more likely. Second, the interview schedule was designed primarily with closed-ended questions. Therefore, the adolescent could respond to questions with a simple "yes" or "no", a number (as in age), the role of a person (e.g., "a neighbor"), or other one word or phrase answer. Therefore, even if someone were listening to the adolescents' answers, they would hear nothing but simple answers such as this. This strategy appears to have been successful since terminated interviews were very low and consistent with rates found with nonsensitive topics and the large majority, over 99% of the adolescents agreed to answer the most sensitive questions (e.g., sexual assault history).

4. Verification of Survey Authenticity:

Interviewers offered respondents the chance to call a toll-free number to SRBI to verify the authenticity of the survey. Interviewers also offered to send a letter before the interview, if the parent/guardian requested one. These follow-up letters were sent to any parent/guardian who requested one, prior to attempting a child interview. Letters explained the sponsorship of the

study, the general survey purpose, the method by which their household had been selected, and the confidentiality of responses. The concerns of most respondents were relieved by this procedure. However, in cases where further information was desired, respondents were given the telephone number of the Co-Principal Investigators at the MUSC National Crime Victims Research and Treatment Center for further verification. If they still had concerns, they were given the telephone number of the project officer at NIJ. Out of nearly 10,000 parent/guardian and adolescent interviews conducted, there were only two calls to the project Co-Principal Investigators and only one to the NIJ project officer. In addition, no human subject incidents were experienced during the entire field period of the NSA.

5. **Incentives:**

As an incentive for participation, adolescent participants received a certificate of participation in the "National Survey of Adolescents" and a check for five dollars as compensation for their time.

6. **Training of Interviewers:**

Interviewers were employees of SRBI who are highly skilled and experienced in conducting this type of sensitive inquiry. The NCVS and SRBI have successfully completed several similar surveys in the past using many of these same interviewers. In addition to the excellent training and experience that these interviewers have received from SRBI, the investigators provided additional training to the interviewers prior to the onset of the field period. The training was specific to the interviews used in this project and focused on the special needs of adolescent respondents.

7. **Rationale for Conducting the Study by Telephone:**

Telephone survey methods offer a valid and efficient method for collecting information from large representative samples of respondents at a relatively low cost with nonsignificant response bias or detection of critical variables of interest as compared to in-person interview approaches (Weeks, Kulka, Lessler, & Whitmore, 1983; Bradburn, 1984). These issues have been looked at specifically in terms of detection of rates of victimization observed using in-person versus telephone interview methods (Catlin & Murray, 1979). Based on objective police report data, no differences in rates of detection of victimization were observed, supporting both the reliability and validity of the telephone method. One recent study (Paulsen, Crowe, Noyes, & Phohl, 1988) compared telephone and in person assessment of DSM-III Axis I disorders, including anxiety disorder, affective disorders, alcoholism, and no mental disorder using a structured diagnostic interview (Paulsen et al., 1988). Kappa's ranging from .69 to .84 were obtained, even with a delay between in-person and telephone methods of 12 to 19 months (Paulsen et al., 1988). The RDD telephone survey method has also been routinely used to complete the Centers for Disease Control Behavioral Risk Factor Surveillance System which assesses risk behaviors within the adult population. Telephone interviewing is also being used with approximately two-thirds of the sample within the National Youth Survey being conducted by Dr. Delbert Elliott (D. Elliott, 1994, personal communication). No differences have been

noted between telephone and in-person interview procedures in assessed rates of delinquent and/or criminal behavior and substance use or abuse

D. **Survey Instruments and Measurement of Key Constructs:**

Two survey instruments were used in the NSA, a parent survey schedule and an adolescent survey schedule, the latter of which is included in Appendix A. The primary purpose of the parent interview was to obtain permission to interview the adolescents. While parents were asked questions about their family history, knowledge of their adolescents' victimization history and current functioning, the main purpose of these questions were to inform the parent of the nature of the study and familiarize them with the types of questions that would be asked of their adolescents. The major focus of the National Survey of Adolescents was obtaining data from adolescents. Thus, the adolescent survey instrument will be described in detail.

Most components of the adolescent survey were selected on the basis of three factors: a) they had been used before in prior telephone interviews, often with adolescents and often by our research team; b) they had good reliability and validity; and c) they measured key constructs in the hypothetical model we wanted to test. Researchers who study the impact of traumatic events in general population samples face a significant test construction problem in that they lack captive samples of patients or college students that are generally available for use in instrument construction and validation. Moreover, funding agencies are reluctant to support extensive pilot work for instrument development. Therefore, it is often impossible to develop psychometric data for revised telephone versions. However, in most cases, there is every reason to believe that well-developed instruments maintain their psychometric properties in revised telephone versions. Our own work has demonstrated that most of these versions maintain internal consistency reliability and construct validity. However, the nature of the constructs measured often means that the state-of-the-art measurement techniques have few known psychometric properties.

This report focuses on data from the following sections of the NSA adolescent survey:

1. **Biographic/Demographic Characteristics:**

The interview collected standard biographic information about respondents, including age, gender, educational achievement, racial status, family income, family composition and structure, and residential location (central city, SMSA remainder, rural).

2. **Family History of Substance Abuse:**

This section utilized questions from our *National Women's Study* telephone survey regarding a family history of substance abuse. Information obtained includes the number of biological parents having substance abuse-related problems, and the types of substance abuse-related social, occupational, or legal problems experienced by biological parents.

3 Victimization History:

This section measures lifetime history of completed rape, other sexual assault, ~~aggravated~~ assault, other physical assault, and witnessing violence in family, school, and neighborhood settings. There is general agreement that sexual assault is the most difficult type of victimization to screen for, and that screening questions must be explicit, capture the full range of sexual assaults (e.g., not just stranger assaults), and permit determination of whether assaults could be legally defined as forcible rape (Kilpatrick, 1983; Koss, 1993; Von, Kilpatrick, Burgess & Hartman, 1991). Following a procedure we used successfully in the *National Women's Study* and the PTSD Field Trial Study, we identified and obtained descriptive information about up to three sexual assaults per respondents: the first sexual assault, the most recent sexual assault, and the worst sexual assault if other than the first or most recent. Descriptive information about each sexual assault included: series or single event, age at onset, frequency, duration, relationship to the perpetrator, extent of physical injuries sustained, did victim think she/he would be killed or seriously injured, did victim ever tell anyone about the assault, if so, who and when, was it reported to police or other authorities, and outcome variables of the social service or criminal justice process. Extensive pilot testing was done to develop the sexual assault screening questions.

Aggravated assault screening questions were identical to those used in the *National Women's Study* and the PTSD Field Trial study. Other physical assault questions are similar to those used by Dembo et al. (1992). As is the case with sexual assault, descriptive information was obtained about the first, most recent, and worst physical assaults. Observation of serious incidents of violence in home, neighborhood, and school settings were measured using some questions from our NIMH-funded Los Angeles Civil Disturbance study (Hanson, Freedy, Kilpatrick, & Saunders, 1993) and others that were developed for this project.

4. Posttraumatic Stress Disorder (PTSD):

The PTSD measure is a modified version of the DIS measure of PTSD using DSM-III-R criteria that we developed and have used in three major telephone survey projects including those which have assessed adolescent respondents. It has been updated to measure PTSD using DSM-IV criteria. This National Women's Study PTSD Module asks respondents if they have ever had a period of a month or more during which they have experienced each PTSD symptom. When symptoms are content specific, respondents are asked to specify the context of that symptom. This method prevents the exclusion of subjects from the PTSD assessment based on the interviewer's judgment of whether or not a particular event meets Criterion A and allows for the assessment of symptom presence in association with a wide variety of events. Information is then gathered about onset age and recurrence of all symptoms. The PTSD Field Trial Study evaluated the degree of reliability between our structured PTSD measure administered by nonclinicians and the Structure Clinical Interview for DSM-III-R (SCID), the "gold standard" of PTSD measures which is administered by clinicians. The *Kappa* coefficient of agreement between the two measures at the diagnostic level was .77 for lifetime PTSD and .71 for current PTSD (PTSD within the past six months). See Resnick, et al., (1993) for a more thorough description of the NWS PTSD Module.

Information was obtained regarding the age of onset and age at most recent occurrence of in order to assess both lifetime and current disorder. Such information permits investigation of the relationship between a history of PTSD and subsequent risk of adolescent victimization and 2) the association between victimization experiences and subsequent risk of the development of PTSD.

5. Substance Use:

The substance use section of the interview was structured to gather the following information. *Has the respondent ever consumed any of the following substances:* a) alcohol, b) marijuana, c) cocaine, d) heroin or other opiates, e) hallucinogens, f) PCP, steroids, g) inhalants or the following prescription drugs used nonmedical. h) tranquilizers, i) barbiturates, j) amphetamines, or k) prescription pain killers. Next, for each substance used, respondents were classified as nonexperimental users if they had ever used the substance four or more times. Next, nonexperimental users were asked several questions about each substance including: a) age of onset, b) frequency of use within the past year, c) recency of use, d) if appropriate, whether they had used the drug IV, and e) if appropriate, if they had driven a car while high or intoxicated from the substance. Information was also obtained about whether substance use ever produced troubles at school, difficulties with friends, criticism by family members, troubles with police, accidents in a car or accidents at home, or health problems. DSM-IV questions were asked that permitted assessment of whether respondents met diagnostic criteria for alcohol or drug dependency or abuse.

6. Substance Dependence and Abuse:

In addition to this information about use of alcohol and drugs, the interview also gathered information that would permit classification into the following diagnostic criteria:

a. *Lifetime Substance Dependence:*

Closed ended questions following DSM-IV criteria (APA, 1994) were asked to determine whether abuse or dependence criteria were met for each type of substance (alcohol, marijuana, and other drugs). Dependence was defined by presence of three or more of the following symptoms: (1) Tolerance, defined by endorsement of either a need for markedly increased amounts of the substance to become intoxicated/high or to have the desired effect or markedly diminished effect with continued use of the same amount of the substance; (2) Withdrawal, manifested by report of two or more reactions including tachycardia, trembling, sleep disturbance, nausea, increased anxiety, seizures, hallucinations, or marked agitation upon suddenly stopping intake to prevent or stop hangover or other withdrawal symptoms; (3) Report of substance use in larger amounts or over a longer period that was intended; (4) Persistent desire or unsuccessful efforts to cut down or stop substance use; (5) Report of spending a lot of time to get or use substance, or to recover from use; (6) Report that specific substance use caused a reduction or elimination of school, work, social, family or recreational activities; (7) Reports of continued use of substance despite the psychological or physical problems that it caused.

b. *Current Substance Dependence:*

Positive if lifetime criteria for substance dependence were not met and the person reported several of these problems within the year prior to interview.

c. *Lifetime Substance Abuse:*

Determined separately for alcohol, marijuana, and other drugs. Abuse was defined as a maladaptive pattern of substance use leading to impairment as evidenced by one or more of the following problems specific to that substance: (1) Report that use ever caused major problems with family, friends, school or work; (2) Report of substance use in hazardous situations including driving a car or boat, swimming, crossing the street in heavy traffic, or other situations in which the individual might get hurt; (3) Arrests or problems with the police because of substance use, including driving while intoxicated, drunk and disorderly, or stealing to obtain drugs; (4) Report of continued use of substance despite problems with family or friends about the substance use, including fights, arguments, or other relationship problems. In addition to meeting these criteria, the individual must not have met lifetime substance dependence criteria for that particular drug or alcohol;

d. *Current Substance Abuse:*

One or more of the problems listed above (a-d) for determination of lifetime abuse occurring within the year prior to the interview and specific to each type of substance. In addition, the individual must not have met lifetime substance dependence criteria for that particular drug or alcohol.

7. Delinquent Behaviors:

Information was obtained regarding the frequency with which respondents had committed Index Offenses as defined within the National Youth Survey. The survey includes assessment of aggravated assault, sexual assault, gang fights, theft of a motor vehicle, theft of items greater than \$50 in value, breaking into a building/motor vehicle, and aggressive behavior towards students, teachers or others. In addition to the past year, frequency of delinquent behaviors, the age at which delinquent involvement began was determined. According to Ageton (1983), the test-retest reliability index for types of offenses reported at a four week retest was .87.

E. Sample Disposition and Participation Rates:

The previously described sampling procedure identified 5,367 eligible households (i.e., households that contained one or more adolescents between the ages of 12 and 17 years old. Out of these 5,367 eligible households:

- 4,836 parents completed interviews (90.1% of eligible households)

- 4,236 parents gave permission for their adolescent to be interviewed (78.9% of eligible households; 87.6% of cases with completed parent interviews)
- 4,023 adolescent interviews were completed (75.0% of eligible households; 83.2% of households with completed parent interviews; 95.0% of households with parental permission).

The recruitment strategy required completing interviews with parents, then getting permission to interview the adolescent, then obtaining permission from the adolescent prior to actually completing the adolescent interview. Given these constraints, we believe that the 75.0% participation and completion rate was quite good.

F. Demographic Characteristics of Parent and Adolescent Samples:

1. Parent Sample:

The sampling strategy permitted the interviewer to interview either parent or guardian in eligible households, so the parent sample (N=4,023) was not a strict probability sample *per se* because parents were not selected randomly within households. Demographic characteristics of the parent sample are presented in Table 1. As inspection of Table 1 reveals, this sample had more women (N=2,887; 71.8%) than men (N=1,136; 28.2%). The majority of respondents were the adolescents' biological parents (90.0%). In 61.1% of the cases, the target adolescent's other biological parent lived in the same household. Over three quarters were married (77.0%), and almost eight out of 10 were employed full-time (79.0%). Slightly more than a third of this sample had annual household incomes greater than \$50,000 (33.9%), more than four out of ten had household incomes between \$20,000 and \$50,000 (43.9%), and nearly 12% had household incomes below \$20,000. With respect to the highest educational achievement attained, 30.2% were college graduates, 59.4% were high school graduates, and 10.4% had less than a high school education.

With respect to ethnicity, 6.7% identified themselves as Hispanic; 92.9% said they were non-Hispanic, and 0.4% refused to answer this question. Non-Hispanic Caucasians were, as expected, the most prevalent racial group (75.9%). Non-Hispanic African-Americans were nearly 15% of the sample. Non-Hispanic, Native Americans and Asians were, in this sample, each less than 1%. Less than 1% of the sample did not give a racial identification.

2. The Adolescent Sample:

In order to better generalize to the U.S. adolescent population, the full sample was weighted to conform to 1995 Census estimates for American adolescents on age, race, and gender. Demographic characteristics for the full weighted sample of N=4,023 are presented in Table 2. The weighted sample included slightly more males (n=2,065) than females (n=1,958) and was composed primarily of non-Hispanic Whites (70.2%). African Americans accounted for nearly 15% of the sample, and Hispanics were nearly 8%. Native Americans were 3.5%, Asians were 1.1%, and Other racial or ethnic identifications were 2% of the sample. Only 28

respondents refused to give or did not know their racial/ethnic identification. The sample was divided fairly evenly across the 12 to 17 age cohorts, with each cohort having approximately 16.7% of the sample. Only 8 respondents refused or did not know their age. Grade level was a bit more diverse, with 8.2% of the sample in the fifth or sixth grades and only 8.2% in the twelfth grade. Other grades ranged from 13.9% to 18.5% of the sample. Only 43 (1.1%) of the respondents did not attend school.

V RESULTS

A Overview of Data Analytic Strategy

Analyses were conducted in four stages. In the first stage, analyses were conducted to provide descriptive information about the prevalence and descriptive characteristics of major study variables. In the second stage, major personal victimization, mental health, substance use, and delinquency variables were broken down by gender, age, and race/ethnicity to permit evaluation of these demographic variables as risk factors. In the third stage of analyses the previously outlined hypotheses were tested. In the fourth stage, follow-up analyses were conducted to investigate interesting findings and/or research questions that had not been hypothesized previously.

B. Descriptive Results

1. Family Problem Substance Use Variables

Overall, 528 adolescents, or 13.1% of the sample, reported that a family member or someone who lived with the adolescent drank alcohol so much they got into fights with other people, beat the children, couldn't get out of bed the next day, or had difficulty holding a job. The prevalence of having a family member with an alcohol problem was significantly related to the adolescent's age. For 12, 13, 14, 15, 16, and 17 years old, respective prevalence rates were 9.3%, 9.9%, 12.4%, 14.1%, 15.4%, and 17.9%. Thus, older adolescents were more likely than young adolescents to report family alcohol problems. Female adolescents were significantly more likely than male adolescents to report family alcohol problems (15.0% vs 11.3%). White adolescents were less likely than nonwhite adolescents to report family alcohol problems (12.2% vs 16.1%).

With respect to having a family member who used hard drugs or had a drug problem, 8.9% of all adolescents (n=358) reported having a family member with such problems. This variable was significantly related to age, with respective prevalence rates for 12, 13, 14, 15, 16, and 17 years old being 5.7%, 7.2%, 9.0%, 11.7%, 11.5%, and 9.4%. Female adolescents were more likely to report such family problems than male adolescents (7.0% vs 10.8%), and nonwhite adolescents were more likely to report than white adolescents (10.9% vs 8.2%).

2. Personal Victimization

a. *Sexual Assault*

(1) *Lifetime Prevalence:*

A total of 326 adolescents, or 8.1% of the sample, reported having experienced at least one sexual assault prior to the interview. Of those who reported having experienced at least one sexual assault, 58.3% had experienced only one, and 41.7% had experienced more than one.

(2) *Age at Time of the Assault:*

As is depicted in Figure 1, the 326 adolescents reporting having ever been sexually assaulted had experienced 462 cases of sexual assault. When asked about their age at the time they were assaulted, 29.9% said they were less than age 11; 16.3% said they were 11 or 12; 20.8% said they were 13 or 14; 20.8% said they were 15 or 16; 1.7% said they were 17, and the remaining 8.7% said they were not sure or refused to answer.

(3) *Relationship to the Perpetrator:*

Presented in Figure 2 is a breakdown of the relationship between the victim and the perpetrator in the 462 sexual assault cases. Almost one third of sexual assault cases (32.5%) involved perpetrators who were friends, and 23.5% of perpetrators were strangers, defined as someone the victims had never seen before or who they had seen before but did not know well. Other types of perpetrators included fathers (1.3%), stepfathers (3.2%), brothers or stepbrothers (1.4%), sisters or stepsisters (0.4%), grandparents (2.8%), other adult relatives (6.5%), other child relatives (6.5%), neighbors (5.8%), coworkers (1.1%), other children (6.9%), or other adults (4.5%). Some adolescents were not sure or refused to answer this questions (3.2%), and these percentages may total more than 100% because some sexual assaults including more than one perpetrator.

(4) *Location of Assault:*

Victims were asked the location in which the sexual assault occurred. As depicted in Figure 3, the most frequent locations were the victims' house (30.5%), the victims neighborhood (23.8%), the victims school (15.4%), or a friend's house (11.8%). Other locations were at a relatives' house (7.8%), outside the victims neighborhood (2.2%), or somewhere else (8%). A few victims (1.3%) refused to answer this question or were not sure about the location.

(5) *Life Threat and Physical Injury:*

Many victims were afraid that they might have been killed or severely injured during the assault. Figure 4 displays their responses. Slightly more than one in four victims (28.1%) said they feared death or serious injury during their sexual assault. In the remaining sexual assault cases, victims said they had no such fears (69.5%) or were not sure or refused to answer the question (2.4%).

With respect to physical injuries, only 1.3% of sexual assault cases resulted in serious injuries, and 11% resulted in minor injuries. The remaining sexual assault cases either produced no physical injuries (85.5%) or a victim who was not sure of the degree of injury or refused to answer (2.2%).

16. *Reporting of Cases to Authorities:*

Victims were asked if their case had ever been reported to police or other authorities. As inspection of Figure 5 indicates, the vast majority of sexual assault cases were never reported to police or other authorities (85.7%). However, 13% of cases were reported to police, 5.8% to child protective services, 5% to school authorities, and 1.3% to other authorities. In four percent of the cases, victims were not sure whether cases were reported or refused to answer this question.

b. *Physical Assault:*

(1) *Lifetime Prevalence:*

A total of 701 adolescents, or 17.4% of the sample, had been victims of at least one physical assault. Of those who had experienced at least one physical assault (50.1%) had experienced one such assault, and 44.9% had experienced more than one.

(2) *Age at Time of Assault:*

Figure 6 presents information on victims; age at the time of the 1054 cases of physical assault. At the time of these physical assault cases, 21.3% of victims were under the age of 11; 21.3% were 11 or 12; 27.1% were 13 or 14; 21.7% were 15 or 16; 4.1% were 17, and 4.2% were not sure or refused to answer this question.

(3) *Relationship to the Perpetrator:*

As Figure 7 indicates, perpetrators were strangers in slightly more than one-third of the cases (36.4%), and 20.5% were identified as friends. Other perpetrators were family members including mothers (3.9%), fathers (4.3%), stepfathers (1.1%), or step-brothers (8.5%), sister or step-sisters (3.9%), grandparents (0.2%), other adult relatives (1.6%) and other child relatives (2.8%). neighbors (4.2%), coworkers (0.2%) were also identified as perpetrators in some cases. Victims were unable to identify the perpetrator or refused to answer in 1.7% of cases.

(4) *Location of Assault:*

Presented in Figure 8 is a breakdown of the location in which physical assault cases occurred. Assaults were most likely to have occurred near the victim's neighborhood (34.2%), in the victim's house (27.9%) and at school (20.2%). Assaults were less likely to occur outside the victim's neighborhood (4.7%), at a friend's house (2.9%), at a relative's house (1.6%) or somewhere else.

(5) *Life Threat and Physical Injury:*

Physical assault victims were asked if they feared being seriously injured or killed during the assault. As depicted in Figure 9, over half of physically assaulted adolescents (52.4%) said

they feared being seriously injured or killed. In the remaining physical assault cases, victims said they had no such fears (44.3%) or were not sure or refused to answer this question (3.2%).

Also included in Figure 9 is a breakdown of the degree of physical injuries sustained in physical assault cases. Almost half of victims reported no physical injuries (47.5%). The remaining cases resulted in minor injuries (45.1%), serious injuries (4.5%) or in the victims not being sure about the extent of injuries or refusing to answer (2.9%).

(6) *Reporting to Police or Other Authorities:*

Presented in Figure 10 is the extent to which physical assault cases were reported to police and or other authorities. As is apparent, 65% of cases were never reported to any authorities, and adolescents in 2.8% of cases were not sure if reports had been made or not, or refused to answer the question. Cases that were reported were reported to police (16.9%), school authorities (16.3%), other authorities (3.8%), and child protection agencies (2.8%).

c. *Physically Abusive Punishment:*

Almost one out of ten adolescents (9.4%) had been victims of at least one incident of physically abusive punishment prior to the interview.

3. Witnessing Violence

Overall, 39.4% of the sample of adolescents reported having witnessed one or more serious incidents of violence. As Figure 11 indicates, 5% of adolescents had seen someone shot with a gun; 10.6% had seen someone cut or stabbed with a knife; 2.8% had witnessed a sexual assault; 10.4% had witnessed a mugging or robbery, and 33.5% had witnessed someone threatened with a weapon.

4. Posttraumatic Stress Disorder

Among this sample of adolescents, 8.1% met DSM-IV diagnostic criteria for PTSD at some time during their lifetime (Lifetime PTSD). A total of 4.9% of these adolescents had PTSD at the time of the interview (Current PTSD).

5. Substance Use/Abuse Dependence

a. *Past-Year and Lifetime Alcohol, Marijuana, and Hard Drug Use:*

The rate of past year alcohol use was 39.8% in the total sample, and 53.9% of the sample indicated that they had used alcohol at least once during their lives (recall that the sample includes children aged 12-17 years). The rates of past-year and lifetime marijuana use were 8.6% and 14.5%, respectively; and the rates of past year and lifetime hard drug use were 2.2% and 9.6%.

b. ***Past-Year and Lifetime Alcohol, Marijuana, and Hard Drug Abuse or Dependence:***

Utilizing DSM-IV diagnostic criteria, the rate of past year alcohol abuse or dependence in the total sample was 3.9%. The lifetime alcohol abuse/dependence rate was 5.6%. Similarly, the rate of past-year marijuana abuse or dependence was 3.7%, with a lifetime abuse or dependence rate of 4.5%. Finally, the rate of past year hard drug abuse or dependence was 0.9%, whereas the rate of lifetime hard drug problem use was 1.2%.

c. ***Past-Year and Lifetime Abuse or Dependence of Any Substance:***

Overall, 6.2% of the sample met DSM-IV diagnostic criteria for past year abuse of a substance (alcohol, marijuana, or hard drugs); 0.7% met criteria for past-year substance dependence and 6.89% met criteria for past year substance abuse or dependence.

6. **Delinquent Behavior**

Among these adolescents, one of eight (12.3%) reported having committed at least one delinquent offense sometime during their life prior to the interview, and nearly one in ten (9.5%) reported having committed at least one delinquent behavior during the year prior to the interview.

C. **Rates of Personal Victimization, Witnessing Violence, PTSD, Substance Use/Abuse/Dependence, and Delinquency Variables by Demographic Characteristics**

1. **Introduction**

One advantage of the NSA sample is that it included large numbers of males and females, different age cohorts, and adolescents of different racial/ethnic groups. This permits analysis on how various demographic groups differ with respect to major variables of interest. Such comparisons also permit testing of several project hypotheses that there will be gender differences in rates of personal victimization, PTSD, substance use/abuse/dependence, and delinquent behavior.

2. **Sexual Assault**

a. ***Gender:***

As inspection of Table 3 indicates, the hypothesis that female adolescents would have significantly higher prevalence rates of sexual assault than males was supported. Overall, the lifetime prevalence of any sexual assault was 13.0% for females and 3.4% for males. Of the types of sexual assault, all were more prevalent among females except oral contact. Only males were asked if they had ever been forced to penetrate others against their will.

b. *Age*

Prevalence of sexual assault was positively associated with age as is depicted in Figure 12. For example, lifetime prevalence rates for 12 and 13 year olds were 3.6% and 4.9%, whereas those for 16 and 17 year olds were 13.7% and 11.9%.

c. *Race/Ethnicity:*

The lifetime prevalence of sexual assault was significantly higher among non-Caucasians than among Caucasians (12.5% vs 6.7%).

3. Physical Assault

a. *Gender:*

As hypothesized, male adolescents had higher prevalence rates of any physical assault than female adolescents (21.3% vs 13.4%). Rates of individual types of physical assault broken down by gender are presented in Table 4. All were more prevalent among boys other than attacked with intent to kill or injure.

b. *Age:*

As was the case for sexual assault, the prevalence of physical assault increased significantly over age cohorts (see Figure 13). It is noteworthy that almost one in four 17 year olds had experienced a physical assault (24.1%).

c. *Race/Ethnicity:*

The lifetime prevalence of physical assault was higher among non-Caucasians than among Caucasians (22.9% vs 15.7%).

4. Physically Abusive Punishment:

a. *Gender:*

As inspection of Table 5 indicates, male and female adolescents did not differ significantly in the prevalence of physically abusive punishment. Females were slightly more likely than males to be spanked so hard they had to see a doctor.

b. *Age:*

Figure 14 presents the lifetime prevalence of physically abusive punishment broken down by age cohorts. Older adolescents were significantly more likely to have experienced physically abusive punishment than younger adolescents.

c. *Race/Ethnicity:*

The lifetime prevalence of physically abusive punishment was significantly higher among non-Caucasians than among Caucasian adolescents (14.3% vs 7.8%).

5. Witnessing Violence

a. *Gender:*

Male adolescents were significantly more likely than female adolescents to have ever witnessed violence (see Table 6). All forms of violence were more likely to be witnessed by males than females *except* witnessing a sexual assault. Girls were twice as likely as boys to have witnessed a sexual assault.

b. *Age:*

As is evident from inspection of Figure 15, the percentage of adolescents who had ever witnessed violence increased significantly over age cohorts. Approximately one in four 12 year olds had witnessed violence (26.9%), but nearly half of 17 year olds had (49.5%)."

c. *Race/Ethnicity:*

Non-Caucasians were significantly more likely to have witnessed violence than Caucasians (54.5% vs 34.7%).

6. PTSD

a. *Gender:*

Female adolescents were significantly more likely than male adolescents to have lifetime PTSD (10.1% vs 6.2%) and to have current PTSD (6.2% vs 3.7%).

b. *Age:*

The prevalence rates of lifetime and current PTSD by age cohorts are presented in Figure 16 and clearly document that rates of PTSD increase significantly with increasing age. It is noteworthy that the rates of lifetime and current PTSD among 17 year olds were 13.1% and 8.4% respectively.

c. *Race/Ethnicity:*

The rate of lifetime PTSD was significantly higher among non-Caucasian than among Caucasian adolescents (9.9% vs 7.3%). Likewise, the rate of current PTSD was significantly higher among non-Caucasian adolescents (6.7% vs 4.1%).

7. Substance Use

a. *Gender.*

There was no significant difference in the percentage of male and female adolescents who had ever used alcohol (54.1% vs 53.7%), although the prevalence of heavy alcohol use during the past year was significantly higher among male than among female adolescents (16.9% vs 13.4%). Rates of lifetime use of illicit drugs non-experimentally did not differ significantly among male and female adolescents (10.2% vs 10.4%). Past year drug use also did not differ as a function of gender (9.4% vs 8.3%).

b. *Age:*

Presented in Figure 17 is a breakdown of rates of lifetime alcohol use and past year heavy alcohol use by age cohort. For both of these variables, there was a significant relationship between age and increased likelihood of alcohol use. For example, almost one in four 12 year olds had used alcohol (24.7%), but almost three-quarters of 17 year olds had (73.9%). Only 2.8% of 12 year olds had past year heavy alcohol use, but almost one-third of 17 years olds reported heavy use of alcohol (31.2%).

A similar pattern emerged for lifetime and past year use of illicit drugs, as is depicted in Figure 18. Whereas only 0.6% of 12 year olds had used illicit drugs, 20.6% of 17 year olds reported such use. Likewise, only 0.4% of 12 year olds reported past year drug use, but 17.8% of 17 year olds did. As depicted in Figure 18, past year illicit drug use closely tracks lifetime use. This trend indicated that most teenagers who begin to use illicit drugs continue to do so.

c. *Race/Ethnicity:*

Lifetime alcohol use was significantly, although only slightly, higher among Caucasian than among non-Caucasian adolescents (55.4% vs 50.5%). Similarly, the proportion of past year heavy alcohol use was also significantly higher among Caucasian adolescents (15.9% vs 13.6%).

There was a small but statistically significant difference in lifetime drug use between Caucasians and non-Caucasians, with the former having higher rates than the latter (10.7% vs 9.4%). The same was true for past year drug use (9.3% vs 7.9%).

8. Current Substance Abuse/Dependence

a. *Gender:*

Male adolescents were significantly more likely than female adolescents to have met DSM-IV diagnostic criteria for alcohol abuse or dependence (4.5% vs 3.2%). They were also significantly more likely to have met diagnostic criteria for marijuana abuse or dependence (4.3% vs 3.0%), but the rates of past year hard drug abuse or dependence were identical for male and female adolescents (0.9% vs 0.9%).

b. *Age:*

reported in Figure 19 are the past year rates of alcohol, marijuana, and hard drug abuse/dependence broken down by age cohort. Each type of abuse/dependence was significantly related to age. These findings reflect the extent to which risk of alcohol and drug problems increase dramatically over the period of adolescence. For example, rates of alcohol, marijuana, and hard drug abuse/dependence among 12 year olds were 0.32%, 0.2%, and 0.0% respectively. Corresponding rates among 17 year olds were 10.4%, 6.9%, and 2.1%.

c. *Race/Ethnicity:*

There were no significant differences between non-whites and whites in terms of alcohol abuse/dependence (3.0% vs. 4.2%, respectively), marijuana abuse/dependence (3.0% vs. 3.9%), or hard drug abuse/dependence (0.7% vs. 1.0%).

9. **Delinquent Behaviors:**

a. *Gender:*

As hypothesized, male adolescents were significantly more likely than female adolescents to have ever committed a delinquent offense (17.7% vs 6.7%) and to have committed one during the past year (13.8% vs 5.0%).

b. *Age:*

The proportion of adolescents who had ever committed an index delinquent offense and who had committed one during the past year increased significantly with age, as is depicted in Figure 20.

c. *Race/Ethnicity:*

Non-Caucasians had significantly higher rates of lifetime index delinquent offenses than did Caucasians (18.1% vs 9.9%). This was also the case for past year delinquent offenses (14.5% vs 7.4%).

D. **Univariate Relationships Between Victimization Variables and Dependent Variables (PTSD, Any Substance Abuse/Dependence, Past-Year Delinquency)**

1. **Posttraumatic Stress Disorder (Current and Lifetime)**

a. *Sexual Assault and PTSD:*

As presented in Figure 21, of male adolescents with one sexual assault, 17.3% met criteria for current PTSD, and 19.4% met lifetime criteria for PTSD; 37.9% of boys who

experienced more than one sexual assault were positive for current PTSD, and 40.6% were positive for lifetime PTSD. Rates of current and lifetime PTSD in boys who had not been sexually assaulted were 2.9% and 5.4%, respectively.

For girls, one sexual assault was associated with a current rate of PTSD of 16.7% and a lifetime rate of 27.0%. Girls experiencing multiple sexual assaults were at greater risk, with 19.5% presenting with current PTSD, and 33.9% having lifetime PTSD. Comparatively, 4.4% of female adolescents with no assaults met criteria for current PTSD and 7.1% met criteria for lifetime PTSD.

b. *Physical Assault or Physically Abusive Punishment and PTSD.*

For male respondents, 6.9% who experienced one instance of physical assault or abusive punishment had current PTSD, and 11.9% had lifetime PTSD. Two episodes of physical assault resulted in current and lifetime rates of PTSD equal to 13.5% and 20.0%, respectively. The current rate of PTSD in boys who had not been physically assaulted or abused was 1.7%, and the lifetime rate was 3.2%.

In female adolescents, the rate of current PTSD associated with one episode of physical assault or abusive punishment was 13.9%, and the rate of lifetime PTSD was 21.3%. In girls with multiple physical assaults, 23.4% had current PTSD, and 39.5% had lifetime PTSD. By contrast, among girls with no physical assault history, 3.6% had current and 6.0% had lifetime PTSD (see Figure 22).

c. *Witnessed Violence and PTSD:*

Approximately 3.4% of male adolescents who witnessed violence presented with current PTSD, and 7.5% reported lifetime PTSD. The rate of current PTSD in boys who had witnessed more than one act of violence was 12.2%, and the lifetime rate of PTSD was 16.8%. Boys who reported never witnessing violence had a current PTSD rate of 1.2% and a lifetime rate of 2.3%.

Girls who witnessed one act of violence had a current PTSD rate of 9.8% and a lifetime rate of 17.2%. Female adolescents who witnessed multiple acts of violence also had higher rates of PTSD (17.4% current; and 27.3% lifetime). Only 2.8% of female adolescents who reported no history of witnessed violence met current criteria for PTSD, and 4.3% met criteria for lifetime PTSD (see Figure 23).

2. Any Substance Abuse/Dependence (Current and Lifetime)

a. *Sexual Assault and any Substance Abuse/Dependence:*

About 27.3% of boys who had been sexually assaulted demonstrated current problematic substance use, and 34.4% had problem substance use during their lifetimes. Rates of current and lifetime substance abuse/dependence in boys who had not been sexually assaulted were 7.1% and 9.0%, respectively.

For girls, sexual assault was associated with a current substance use abuse rate of 27.5% and a lifetime rate of 27.5%. Comparatively, 3.6% of female adolescents who had not been sexually assaulted were current substance abusers and 5.4% were lifetime substance abusers (see Figure 24)

b. *Physical Assault and any Substance Abuse/Dependence:*

Rates of problematic substance use in boys following physical assault were also high, with 19.4% of male respondents reporting current substance abuse/dependence and 24.0% reporting lifetime abuse/dependence. Only 4.7% of boys who were not physically assaulted were current substance abusers, and 6.1% were lifetime abusers.

Rates for girls mirrored those of boys. Physical assault was associated with a current substance abuse/dependence rate of 20.1% and a lifetime rate of 26.4% in female adolescents. Girls who were not physically assaulted had lower rates of current and lifetime use (3.7% and 5.5%, respectively) (see Figure 24).

c. *Physically Abusive Punishment and any Substance Abuse/Dependence:*

This form of punishment produced rates of problematic substance use similar to those produced by physical assault in boys, but produced somewhat less pronounced effects in girls. Specifically, 19.7% of boys who experienced physically abusive punishment reported current substance abuse/dependence, and 23.4% reported lifetime problematic substance use. By contrast, 6.7% of boys who were not abused reported current problematic substance use, and 8.6% reported problematic substance use at some time during their life.

Approximately 12.4% of female adolescents who were physically abused reported that they were currently substance abusers/dependent, and 17.3% were lifetime problem substance users. Five percent of girls who did not experience physically abusive punishment were current substance abusers and 7.2% reported lifetime substance abuse/dependence (see Figure 24).

d. *Witnessed Violence and any Substance Abuse/Dependence:*

Approximately 13.9% of boys who witnessed violence were current substance abusers, and 17.0% were lifetime abusers. However, only 3.2% of boys who did not witness violence reported current substance abuse/dependence, and 4.4% reported lifetime problematic substance use.

Rates for girls again paralleled those of boys. About 13.2% of female adolescents who had witnessed violence were current substance abusers/dependent, and 17.8% had lifetime substance abuse/dependence. By contrast, only 2.0% of girls who did not witness violence evinced current, and 3.1% evinced lifetime substance abuser or dependence (see Figure 24).

Past Year Delinquency

Sexual Assault and Past Year Delinquency

Fully 41% of the sexually assaulted males reported engaging in delinquent acts, compared to 13% of those not sexually assaulted. Fewer (15%) sexually assaulted females reported engaging in delinquent acts, but this rate was five times higher than that for girls who were not sexually assaulted (3%).

b Physical Assault and Past Year Delinquency:

The proportion of physically assaulted boys who engaged in past year delinquent acts was 36.9%, compared to 7.5% of non-assaulted boys. Similarly, 23.5% of physically assaulted female adolescents reported past year delinquent acts compared to 2.1% of non-assaulted girls (see Figure 25).

c. Physically Abusive Punishment and Past Year Delinquency:

The results for physically abusive punishment resembled those of assault in that 35.8% of abused boys engaged in delinquent acts compared to 11.7% of non-abused boys. Approximately 15.4% of abused girls participated in past-year delinquent activities, relative to 3.8% of non-abused girls.

d Witnessed Violence and Past Year Delinquency:

About a fourth (25%) of males who witnessed violence reported engaging in delinquent acts, compared to only 5% of boys who did not witness violence. About 13% of girls who witnessed violence reported delinquency, compared to 1% of girls who did not witness violence.

E. Multivariate Analyses of the Relationship Between Violence Exposure and Posttraumatic Stress Disorder

1. Overview of Analyses

The primary objective of this set of analyses was to test the hypothesis that multiple exposure to sexual assault, physical assault, and witnessing violent events increases risk of PTSD after controlling for the effects of other variables that might be expected to influence risk of PTSD such as demographic characteristics and family environment. As was previously noted, many adolescents had experienced more than one incident of violence, so we constructed the following measures of multiple exposure to violent incidents using previously-described variables:

- Number of Sexual Assaults Exposures:
Score: 0=none; 1=one; 2=more than one

- Number of Physical Assault Exposures:
Note: this included physical assaults and physically abusive punishment
Score: 0=none; 1=one; 2=more than one
- Number of Witness to Violence Exposures:
Score: 0=none; 1=one; 2=more than one
- Total Violence Exposure: The sum of sexual assault, physical assault and witnessing violence scores.
Scores: 0= no exposure; 6=more than one exposure to each of the three types of violence.

Three types of data analyses were conducted:

- Rates of violence exposure were compared by gender
- Analyses were conducted to determine whether demographic, family environment, and violence exposure variables were risk factors for lifetime PTSD.
- Multivariate logistic regression analyses were used to test the hypothesis that extent of multiple exposure to violence increases risk of PTSD after controlling for the effects of demographic and family violence variables.

2. Results of Univariate Analyses

As has been described previously, rates of each type of violence differed significantly by gender as well as by race in most cases. Presented in Figure 28 are results of analyses comparing rates of PTSD among male and female adolescents with and without family members with alcohol or drug problems. As inspection of this figure illustrates, female adolescents had higher rates of PTSD than male adolescents, and adolescents with family members who had alcohol or drug problems had higher rates of PTSD than those who did not. Figure 27 provides information about the number of violent incidents that male and female adolescents had experienced or witnessed.

3. Results of Multivariate Logistic Regression Analysis

To test the hypothesis that extent of multiple exposure to violence would increase risk of PTSD after controlling for the effects of other variables, a hierarchical multivariate logistic regression analysis was conducted using lifetime PTSD as the dependent variable. In Step One of this analysis, the demographic variables age, gender, race, and income were entered simultaneously. In the second step family alcohol problems and family drug problems were entered. In the final step, the number of sexual assaults, physical assaults, and number of incidents of violence witnessed were entered.

Table 7 includes the results of this analysis. Odds ratio in the Step OR column represent the increase in odds of PTSD controlling for the effect of other variables in the model either

entered at the step and all variables entered at earlier steps. Odds ratio in the Final OR column represent the unique increase in odds of PTSD when controlling for the effects of all other variables in the final model after all variables have been entered.

At Step 1, all demographic variables significantly increased odds of PTSD after controlling for the effects of other demographic variables. Increased age was associated with a substantial increase in odds of PTSD. The age variable had six values (i.e., 12, 13, 14, 15, 16, 17 years old), so there were five potential steps of increase in age. For such multilevel variables, the states OR represents the increase in odds associated with a one level difference. Thus, the odds increase between 12 and 13 year olds or between 16 and 17 years in PTSD likelihood was 1.28. However, the odds increase at a two step difference is the OR squared; for a three step difference, it is the OR cubed, etc. Thus, the increase in PTSD odds is 1.28^2 , or 3.44. Odds of PTSD were 1.69 times higher for female than for male adolescents. Odds of PTSD were 1.34 times higher for nonwhite adolescents than for white adolescents. PTSD was not related to household income.

In Step 2 of the analyses, the family alcohol problem and family drug problem variables were entered. Both of these variables were significantly related to odds of PTSD after controlling for the effects for each other as well as for the demographic variables in Step 1. Odds of PTSD were 2.80 times greater among adolescents with a family alcohol problem. Likewise, odds of PTSD were 2.36 times greater among adolescents with a family drug problem than among those without a family drug problem.

In Step 3 of the analyses, three variables measuring the number of sexual assaults (i.e., 0,1, or more than one), physical assaults (i.e., 0,1, or more than one), and incidents of violence witnessed (0,1, or more than one) were entered. The resulting OR represented the unique increase in odds of PTSD controlling for the effects of the other two variables previously entered at the first two steps. As the results of these analyses indicate, each of the three exposure to violence variables increased odds of PTSD significantly after controlling for each other and all other variables in the model. Odds of PTSD were 1.69 times higher among victims of one sexual assault vs. adolescents with no sexual assault. Compared to adolescents without a sexual assault, victims of more than one sexual assault had odds 2.86 times greater of PTSD. Compared to adolescents with no physical assaults, odds of PTSD were 1.75 times greater among adolescents with one physical assault and 3.06 times greater for those who had two or more assaults. Adolescents who had witnessed one incident of violence were 1.91 times greater to have PTSD, and those who witnessed more than one incident of violence had odds 3.69 times greater.

Inspection of the final odds ratio in Table 7 reveals several interesting findings. First, the effects of the demographic variable of race became nonsignificant after controlling for the effects of family alcohol and drug problems and the number of sexual assaults, physically assaults, and violence incidents witnessed. Second, the effects of age and gender remained significant but effects of age were reduced in magnitude, suggesting that its effect on odds of PTSD were at least partially mediated by the effect of family alcohol or drug problems and exposure to violence. Third, the effects of family alcohol or drug problems also remained significant but were reduced in magnitude after controlling for violence exposure, suggesting that the effects of these variables too appear to be at least partially mediated by violence exposure.

These findings clearly supported the hypothesis that extent of exposure to violence is an important risk factor for PTSD even after controlling for a host of other relevant variables. This finding is clearly illustrated in Figures 21-23, which depict rates of lifetime PTSD as a function of the number of violent incidents adolescents had experienced or witnessed.

F. Multivariate Analyses of the Relationship Between Violence Exposure and Past Year Substance Abuse/Dependence

1. Overview of Analyses

The major objective of this set of analyses was to test key project hypothesis about the relationships between exposure to violence and likelihood of developing substance abuse and dependence. The first hypothesis is that exposure to violence will increase likelihood of substance abuse and dependence. A second hypothesis is that PTSD will also increase risk of substance abuse and dependence. The final hypothesis is that exposure to violence and developing PTSD will increase risk of substance abuse/dependence after controlling for relevant demographic and family history variables.

These analyses focused on substance abuse/dependence rather than substance use. Adolescents who meet DSM-IV diagnostic criteria for substance abuse or dependence are clearly experiencing major problems associated with their use of alcohol or other drugs, and it is important to understand factors associated with those adolescents who are having the most problems associated with their substance use.

Analyses were conducted separately for past year alcohol abuse/dependence, past year marijuana abuse/dependence, and hard drug abuse/dependence. DSM IV criteria were used to determine whether each adolescent had met criteria for alcohol, marijuana, or hard drug abuse or dependence within the past year.

Two types of analyses were conducted:

- Univariate analyses compared the risk of alcohol abuse/dependence, marijuana abuse/dependence, and hard drug abuse/dependence as a function of demographic variables, family history variables, and PTSD.
- Multivariate logistic regression analyses tested the hypothesis that exposure to violence and having current PTSD will increase risk of alcohol, marijuana, and hard drug abuse/dependence after controlling for the effects of demographic and family history variables.

2. Results of Univariate Analyses

a. *Alcohol Abuse/Dependence*

As is depicted in Figure 28, victims of sexual assault, physical assault, and those who had witnessed violence had higher rates of alcohol abuse/dependence than their counterparts who had never been exposed to these types of violence. Likewise, adolescents with family alcohol or drug problems also had higher rates of alcohol abuse/dependence than those who did not. Rates of alcohol abuse/dependence were also higher among those adolescents who had current PTSD than among those without current PTSD. The results of these analyses, as well as those examining demographic variables as risk factors for alcohol abuse/dependence, are presented in Table 8, section A. With respect to these demographic variables, age and income were significantly related to odds of alcohol abuse/dependence. Gender was related but not significantly so after the Bonferroni correction, which required $p < .01$ to maintain setwise alpha at .096.

b. *Marijuana Abuse/Dependence*

A similar analysis using marijuana abuse/dependence as the dependent variable is presented in Figure 29 and in the second section of Table 8, section B. As was the case with alcohol abuse/dependence, victims of sexual assault, physical assault or physically abusive punishment and witnessing violence were much more likely to have marijuana abuse/dependence than adolescents who were not exposed to violence. Adolescents with family alcohol or drug problems and those with current PTSD were also at higher risk for marijuana abuse/dependence than their counterparts without such family histories or PTSD. Age was a risk factor for marijuana abuse/dependence, but gender, race, and income were not.

c. *Hard Drug Abuse/Dependence*

The analysis of risk for hard drug abuse/dependence is depicted in Figure 30 and in the third section of Table 8, section C. A similar pattern of results emerged, with history of sexual assault, physical assault or abusive punishment, and witnessing violence significantly increasing odds of hard drug abuse/dependence. The same was true for family alcohol or drug problems and for current PTSD. Age was the only demographic variable to increase odds of hard drug abuse/dependence.

3. Multivariate Logistic Regression Analyses

a. *Alcohol Abuse/Dependence:*

Section A of Table 9 includes the results of the multivariable logistic regression analyses examining risk of alcohol abuse/dependence. In the first step of this analysis, the variables of age, gender, race, and income were entered simultaneously. As is apparent, the demographic variables of age, gender, and income all significantly increased odds of alcohol abuse/dependence. When family alcohol and family drug problems were entered in the next step

of the analysis, family alcohol problems increased odds significantly but family drug problems did not. During the third step, sexual assault and physical assault/physical abuse/sexual abuse were entered, and each of these variables increased odds of alcohol abuse/dependence after controlling for the effects of each other and of all other variables that had been previously entered. Witnessing violence was entered in the fourth step and increased odds of PTSD after controlling for all other previously entered variables. When current PTSD was entered in the last step, it did not increase odds of alcohol abuse/dependence after controlling for the effects of all other variables.

Inspection of the OR's in the final model reveals several interesting findings. First, when the effects of all other variables were controlled, the demographic variables of age, gender, race, and income all had a significant relationship to alcohol abuse/dependence. Odds of alcohol abuse/dependence increased substantially with age, as might be expected. Odds were higher among male adolescents than among female adolescents. Odds were higher among Caucasians than among non-Caucasians and were higher among adolescents from less affluent households. Second, the effect of family alcohol problems remained significant after controlling for the effects of all other variables, although odds were reduced in magnitude. Third, exposure to violence increased odds of alcohol abuse/dependence substantially and continued to do so after controlling for other variables, as had been hypothesized. Fourth, current PTSD did not significantly increase odds of alcohol abuse/dependence after controlling for the effects of all other variables as was hypothesized.

b. *Marijuana Abuse/Dependence*

Depicted in section B of Table 9 are the results of the multivariate logistic regression analyses examining odds of marijuana abuse/dependence. In step one, demographic variables were entered simultaneously. Age and male gender increased odds significantly whereas race and income did not. In step two, family alcohol problems and family drug problems increased odds significantly. In step three, adolescents who were sexually assaulted and those who were physically assaulted had significantly higher odds of marijuana abuse/dependence than their counterparts who had never been assaulted. In step four, having witnessed violence significantly increased odds of marijuana abuse/dependence after controlling for effects of all other previously entered variables. In the final step, odds increase significantly among adolescents with current PTSD vs. those who did not after controlling for the effects of all other variables.

In the final model, all demographic variables were significant, with age, male gender, Caucasian race, and lower household income being associated with higher odds of marijuana abuse/dependence. Family drug problems, but not family alcohol problems, also increased odds of marijuana abuse/dependence. As hypothesized, exposure to violence significantly increased odds of marijuana abuse/dependence after controlling for the effects of demographic variables and family drug problems. Also as hypothesized, current PTSD increased odds of marijuana abuse/dependence after controlling for the effects of demographic variables, family drug problems and exposure to violence.

Hard Drug Abuse/Dependence

As described in Section 4 of Table 9, demographic variables of age and income were found to increase odds of hard drug abuse/dependence significantly, but gender and race did not. Family alcohol problems and family drug abuse problems, which were entered in step two, both significantly increased odds of hard drug abuse/dependence. In step 3, sexual assault and physical assault/physically abusive punishment both significantly increased odds of hard drug abuse/dependence. In step 4, witnessing violence increased odds of hard drug abuse/dependence. In the final step, current PTSD was entered but did not significantly increase odds of hard drug abuse/dependence.

In the final model, age and Caucasian race were the demographic variables that substantially increased odds of hard drug abuse/dependence. Both family alcohol and family drug problems significantly increased odds. As hypothesized, exposure to violence increased odds of hard drug abuse/dependence after controlling for the effects of demographic and family alcohol and drug problems. However, the hypothesis that current PTSD would increase odds after controlling for effects of all other variables was not supported.

4. Timing of Exposure to Violence and First Use of Alcohol or Drug

Given the aforementioned strong relationships between exposure to violence and odds of alcohol abuse/dependence, marijuana abuse/dependence, and hard drug abuse/dependence, an obvious question arises as to whether use of substances preceded the violent assaults or *visa versa*. To examine this timing issue, we examined data from 318 adolescents who reported having been a victim of physical or sexual assault and who reported having ever used alcohol, marijuana, or hard drugs. Based on information they provided about the year they were first assaulted and the year they first consumed alcohol, marijuana, and hard drugs, they were classified into three groups for each substance:

- First substance use preceded year of first assault
- First assault preceded year of first substance use.
- ▶ First substance use and first assault occurred during the same year

Next, the proportion of the 318 adolescents falling into each of the three groups for each of the three substances was determined. Only a minority of adolescent victims who had ever used substances said they had used them in a year prior to the year they were assaulted (25.6% for alcohol use, 31.0% for marijuana use, and 19.8% for hard drug use). Other adolescents said that their first substance use and first assault occurred during the same year (20.6% for alcohol, 21.2% for marijuana, and 16.7% for hard drugs). However, the bulk of adolescent victims said that their first use of substances occurred in a year that their first use of substance occurred *after* the year they were first assaulted (53.8% for alcohol, 47.8% for marijuana, and 63.5% for hard drugs).

Unfortunately, it was not possible to determine whether the assault or the substance use occurred within a given year. Also, information was not asked about the year in which incidents of witnessed violence or physically abusive punishment occurred, thereby precluding an analysis of the timing of these events *vis a vis* first use of substances. Nevertheless, it appears that more adolescent assault victims were assaulted before they ever used alcohol or drugs than *visa versa*.

G. Multivariate Analyses of the Relationship Between Violence Exposure and Past Year Delinquency

1. Multivariate analysis of delinquency correlates

Initial multivariate models were prepared for predicting the commission of at least one delinquent (index) offense within the year prior to the survey by dividing the sample by gender. As reported above, the prevalence of past year delinquent acts was higher for males than females. The prevalence of sexual assault was larger for females than males but the prevalence of physical assault was larger for males. These univariate analyses suggest that victimization may play different roles in the development of delinquency behavior for males and females. Therefore, initial models were developed for each gender.

Table 10 presents results from two hierarchical logistical regressions predicting the commission of at least one index offense in the past year, one for male adolescents and one for females. Demographic variables were entered in step 1, family history of substance use variables in step 2, personal victimization variables in step 3, witnessing violence in step 4, lifetime PTSD in step 5 and problem substance use in step 6. Odds ratios for each variable at the time of entry into the model (step) and for the final model are presented.

For male adolescents, when other substantive variables were entered in the model, all the demographic variables were nonsignificant. This finding suggests that while univariate analysis may suggest higher prevalence rates of delinquency in low-income and minority populations, when the other variables in this model are entered, these demographic trends fall out. Physical abuse or assault was the most prominent predictor in this analysis with a step odds ratio of 4.84 and a final model odds ratio of 3.06. These findings indicate that adolescent males with a history of physical assault or physical abuse were three to four times more likely to commit delinquent acts in the past year, even controlling for other significant risk factors such as demographic characteristics, family problem substance use history, and personal problem substance use. In the final model, history of physical abuse and assault was the most prominent predictor other than problem substance use. Witnessing community violence was also a substantial predictor, with adolescent males witnessing violence being three times more likely to commit delinquent acts. Somewhat surprisingly, sexual assault history, while significant in the step entry, was not significant in the final model. History of PTSD, however, was not significant at the step entry, but was significant in the final model. This finding suggests a suppressor effect with problem substance use for males. That is, when the effects of problem substance use are controlled for, PTSD becomes a significant predictor of delinquency for adolescent males. Not surprisingly, family history of problem substance use remained in the final model as significant predictors.

Similar results were found in the predictive model for adolescent females. History of physical assault and abuse were even greater predictors for girls than boys, with an increase in odds of nearly four times. Witnessing community violence also was a stronger predictor among girls than boys, nearly four times increase vs. three for boys. Like boys, substance abuse was the largest predictor of all. Also similar to boys, a history of sexual abuse was a significant predictor at the step it was entered. However, when the final three variables were entered, it was no longer significant. Having a history of PTSD significantly increased the odds for delinquency for girls as well as boys, as hypothesized. Family history of substance abuse was also significant, particularly family problems with drug abuse. While the odds for boys with this family history were increased nearly two times, for girls it was an even stronger predictor, increasing odds nearly three times greater.

In an effort to further specify predictive models with these variables, separate logistic regressions were conducted on individual gender by racial/ethnic identification groups. Unfortunately, even with the relatively large sample size of this study, there were sufficient numbers within individual cells to examine only the White and African-American adolescents in the sample using this approach. Results for White and African-American males are presented in the top half of Table 11, and results for White and African-American females are presented in the bottom half of Table 11. These analyses are slightly different from those above. Only Income was entered as a demographic control since gender and race/ethnic identification were already controlled by limiting and dividing the samples. History of physical assault and history of physical abuse (i.e., harsh physical discipline) were divided into two separate variables for more precise analysis. These four analyses give gender and racial/ethnic group-specific models for predicting past year delinquent behavior.

The differences between the four analyses are the most interesting part of this analysis. Income was only significant for White males, and only at the initial step entry. In all final models it was not significant for any of the groups. This finding indicates that when the other variables are the important factors in predicting delinquency and that family income is not important when these other factors are considered.

Family history of alcohol abuse was significant for all four groups at the step entry, i.e., controlling for income. It remained a significant factor in the final model (i.e., controlling for all other variables) for all groups except African-American males. For this group, it had the smallest step OR, and was not a significant factor in the final model. For white males, family history of alcohol abuse increased the odds of delinquency nearly two times. For both female groups it increased the odds approximately three times. Therefore, African-American males were different from the other gender-racial/ethnic groups. Similar results were found for family history of drug abuse. Step OR's were significant for all groups other than African-American males, and this factor remained significant for all but White females.

Witnessing violence was a strong predictor for all four groups at both the step and final analyses. Interestingly, the size of the effect was smaller for White males compared to the other three groups, though still substantial. The effect size was particularly large for African-American females.

Most surprisingly, history of sexual assault was not a significant predictor for any group either at the step entry or in the final model. This finding was counter to our hypothesis. However, history of physical assault was a significant predictor for all the groups at the step entry. It was also significant in the final model for all groups other than the African-American males.

Having a history of physical abuse (e.g., harsh physical discipline) was a significant predictor only for White males. For this group it was a strong predictor, doubling the odds of delinquent acts. However, it was not significant for any of the other gender-racial-ethnic groups at either the step entry point or in the final model.

Having a lifetime history of Posttraumatic Stress Disorder was a significant predictor only for African-American males. It was significant at the step entry point for White females, but dropped out in the final model for this group. Therefore, again, African-American males predictive models for African-American males appear to differ from the other groups.

As expected, significant problems with substance use (not necessarily abuse or dependence) was related to delinquency behavior, even after controlling for all other factors in the model. This finding was true for all groups at both the step entry points and the final models.

Several conclusions grow out of these analyses. First, it is clear that the pathways to delinquency, at least based on the factors assessed here, may be quite different for boys and girls from different racial and ethnic groups. Specifically, African-American males seem to have some unique issues and less in common with White males or African-American females than might be expected, considering gender and race/ethnic identification individually. Several of the significant predictors for these other groups were not important to African-American males. The family history of problems with substance use variables and history of physical assault or physical abuse were not associated with delinquency with this group as they were in the other groups. Also, a history of PTSD was important for African-American males, but not for the other group. Therefore, these adolescents may have a unique set of circumstances that lead to delinquency. This suggests that future research and prevention efforts should not only take into account gender and race as factors, but should also examine the interaction between these factors.

If one assumes that adolescents are more likely to see violence if they live in more violent communities, this analysis confirms the importance of violence in the community to contributing to delinquency. This factor was a consistent correlate across all of the groups. The same can be said of problems with substance abuse. Clearly, witnessing violence in the community and substance abuse are two significant and substantial correlates of delinquency, regardless of gender or race/ethnic identification.

However, even controlling for these factors, experiencing a physical assault is also a strong correlate, except for African-American males. Therefore, the hypothesized relationship between this form of victimization and delinquency was confirmed. One of the most surprising findings of the analysis was that having a history of sexual assault was not associated with delinquency in any group, controlling for these other variables. This finding was counter to our

implications. It also was surprising that having a history of physical abuse was predictive only for White males. Again, these findings illustrate the importance of specific gender and racial-ethnic

Tables 12 and 13 attempt to put some of these findings into an epidemiological perspective. While correlation is important, and strong multivariate correlations such as were found in the above analyses do enlighten, they do not tell the full story. In these tables, the primary predictor factors were combined by class. That is, the two family history of substance use problems variables were collapsed into one, such that if a participant was positive on either of the two variables, they were positive on the collapsed variable. Similarly, if a participant had a history of sexual assault, physical assault, or physical abuse, they were labeled assault positive. PTSD and problems with substance abuse were the same as the above analyses.

These tables divide the sample into 16 mutually exclusive groups that represent all possible combination of the collapsed predictor variables. For example, the first group is negative for all the risk factors and the last group is positive for all the risk factors. The middle groups are the various combinations of positive and negative risk factors. Two tables are presented, one for male adolescents and one for females.

The far right column of these tables is the percentage of adolescents in that risk factor group (i.e., those adolescents with the indicated set of risk factors) who committed at least one index offense in the year prior to the survey. In Table 13 for adolescent males, in the first group, the group with none of the risk factors, only 4.5% of the adolescent males committed an index offense in the past year. However, in the last group, the group with all the risk factors, nearly 90% of the adolescent males had committed an offense. This appears to be excellent predictive power. In other words, if we locate a male adolescent with all of the risk factors, then he has a 90% chance of having recently committed a delinquent offense. Less impressive, but consistent results were found for adolescent females. In the completely negative risk group, less than 1% of the girls had committed a delinquent offense in the past year. However, in the all positive risk factors group, 43% had done so. The smaller percentages compared to males are due to the fact that the girls had an overall delinquency rate about one-half that of boys. These results seem to have great value for targeting prevention programs and possibly for the development of

However, this conclusion, while accurate, may be misleading. The first column on the right lists the percentage of the adolescents who had committed a delinquent act in the past year who are in each risk factor group. For example, while 90% of the males in the all positive risk factor group had committed a past year delinquent offense, they represented only 6.5% of all the male delinquents. On the other hand, only 4.5% of the male adolescents in the all negative risk factor group had committed a past year delinquent act. But, they represented 20% of all the male delinquents. This apparent paradox is explained by the middle column on the right. This column lists the percentage of the entire male adolescent sample that each risk factor group represents. It can be seen in this column that the all negative risk factor group accounts for 61% of all male adolescents. Therefore, though only 4.5% of the male adolescents in this group had committed an index offense in the past year, because they are 61% of the male adolescent population, they account for 20% of the male delinquents. Similarly, while the all positive risk factor group had a

90% delinquency rate, they represented only 6.5% of all male delinquents because they are only 1% of the male population.

Similar results were found for girls. The no risk factor group had a less than 1% delinquency rate, but they accounted for nearly 9% of the female delinquent youth because they are nearly 60% of the adolescent female population. The all positive risk factor group had a 43% delinquency rate, but accounted for only 12% of the female delinquents, because this group was only 1.3% of the female adolescent populations.

These results illustrate why it is important to understand the epidemiological contour of risk factor analysis and the importance of comparison groups. While a particular risk factor may be strongly related to delinquent behavior, it may be so rare in the general population that it may actually be present in a very small number of delinquents. Understanding the prevalence of risk factors in the general and the delinquent population, therefore, is crucial.

H. Prevalence Summary and Population Estimates of Critical Study Variables

Table 14 summarizes prevalence rates for critical study variables. In addition, this table provides census-based estimates of the number of affected adolescents in the United States on each study parameter. Such "actual number affected" estimates are a noted benefit of the census-driven RDD methodology employed by this study. As is illustrated in Table 14, rates of interpersonal violence and victimization are extremely high in American youth. Almost 2,000,000 (8.1%) children have been sexually assaulted in this country. Fully 3,900,000 (17.4%) have been severely physically assaulted, and another 2,100,000 (9.4%) have been punished in a manner considered physically abusive. Most pervasive is witnessed violence, with approximately 8,800,000 (39.4%) children indicating that they have seen someone shot, stabbed, sexually assaulted, physically assaulted, or threatened with a weapon.

Clearly, victimization of youth in this country is widespread and demands attention. This point is firmly underscored when one considers the emotional impact of rampant assaultive violence on our children. Our population-based estimates indicate that fully 1,800,000 (8.1%) children have met criteria at one point in their lives for lifetime PTSD, and 1,100,000 (4.9%) currently suffer from the disorder. Potentially more damaging is substance abuse that frequently follows assault. Two million (9.1%) youth have met criteria for substance abuse or dependence (i.e., abuse of or dependence on alcohol, marijuana, or hard drugs) at some point in their lives. Approximately 1,500,000 children and adolescents currently are dependent on or abusing substances.

Delinquency estimates are also provided in Table 14. Approximately 2,700,000 (12.3%) youth have committed an delinquent act at some point in their lives, and 2,100,000 (9.5%) have committed such an act in the past year.

These numbers are independently disconcerting, but become even more distressing when one considers the high rates of victimization of American youth and the apparent relationship between victimization and substance abuse or delinquency. That is, victimization, in addition to causing emotional problems such as PTSD and depression, may also lead to substance abuse and other illegal or destructive activities in youth, and ultimately in adulthood. Although the present

cross sectional study clearly established the relationship between victimization and negative outcomes (PTSD, substance abuse, delinquency) longitudinal study designs are required to more firmly establish such causal conclusions.

VI. DISCUSSION

The National Survey of Adolescents demonstrated the feasibility of conducting sensitive, clinically-relevant, large-scale research with this age group using household probability sampling methods and parental consent. Our findings regarding prevalence of sexual and physical assault were consistent with those of other studies in this area (e.g., Saunders et al., 1992). As hypothesized, the rate of sexual assault was higher in females (13%) than males (3.4%), whereas the rate of physical assault was higher in males (21.3%) than females (13.4%). Rates of physically abusive punishment did not differ significantly between males (8.5%) and females (10.2%). Levels of physical assault and sexual assault increased substantially with age, and overall prevalence rates were higher for Native Americans, African Americans, and Hispanics than among Whites. However, rates of assaultive violence were inversely related to income, suggesting that socioeconomic status may mediate differences across ethnicity. Because proximity to, as well as experience of violence may increase risk of negative outcomes, the NSA also included a measure of Witnessed Violence. For all studied variables, including PTSD, substance abuse, and delinquency, having witnessed violence greatly increased risk for male and female adolescents. Approximately 44% of male participants and 35% of female participants reported that they had witnessed a violent act, with 38% of males and 29% of females indicating that they had actually seen someone threatened with a weapon. The NSA also studied prevalence of PTSD in sample. Eight percent of adolescents surveyed met criteria for the disorder, with rates of girls (10%) higher than those for boys (6.2%). The NSA also examined rates of delinquency in American adolescents. Boys were far more likely to have engaged in delinquent acts at some point in their lives than girls (lifetime prevalence: 17.7% males, 6.7% females).

Overall, rates of substance use reported in our sample are lower than those reported in the Monitoring the Future Study (1995) and slightly lower than those reported in the National Household Survey of Drug Abuse (1995), despite the fact that usage queries were very similar in each investigation. Several factors might explain these discrepancies. Foremost among these is methodological variance across studies. While the Monitoring the Future and National Household Survey studies allow adolescents to indicate use on self-completed questionnaires, the present study required adolescents to verbally report use. This might have contributed to lower rates for two reasons. First, respondents might have been reluctant to overtly and personally describe their use of illicit substances to another individual. Second, respondents might have feared describing their patterns of substance use aloud in their parents' home. Slightly lower rates of use do not diminish findings, however. By contrast, the findings related to victimization and substance use are even more robust given the conservative estimates of use prevalence.

Data regarding order of onset of substance abuse and victimization were clearly interpretable and consistent across all classes of drugs. For a large proportion of children, victimization preceded substance use. Relatedly, the important etiological role of victimization (discussed below) in delinquency is further highlighted by this finding.

In order to examine the unique impact of each variable on use and problem use (defined as either substance abuse or dependence) of each substance over and above effects of other variables, five-step hierarchical logistic regression was employed in which odds ratios of variables

entered on each step were adjusted for those of other variables entered both on that step and on preceding steps. Grouping was rationally driven so as to maximize relevance of findings. This was accomplished by adopting a purposefully conservative analytic approach. As such, demographic variables were entered first (i.e., age, race, gender, and income). Effects of familial alcohol and drug use on adolescent substance use were examined in the second step. Of course, any effects noted for these variables would therefore be apparent over and above impact attributable to demographic variables. Victimization variables were entered following demographic and familial variables, to permit strengthening (or weakening) of conclusions that effects of abuse on substance use were not simply the result of uncontrolled sources of correlational variance. Witnessed violence is qualitatively different than experienced violence and was therefore entered on a separate step. PTSD status was entered on the fifth and final step because this variable referred to a diagnostic constellation of symptoms rather than to a specific event or situation. Thus, our interest was whether or not such a constellation of symptoms increased risk of substance use and abuse independent of victimization and familial behavior. Also provided were Final Model odds ratios, which illustrated the unique impact of each variable on substance use while simultaneously controlling for effects of every other variable.

With regard to demographic factors, Caucasian and male respondents were slightly more likely to report alcohol use and problem alcohol use, but this finding was not consistently observed. Older children, and children from higher SES backgrounds were also more likely to engage in problem alcohol use. Risk of *problem use* was nearly tripled in individuals with familial alcohol use, past physical assault/abuse, or sexual assault. Moreover, this increased risk was apparent for these variables even when effects of all other variables were controlled. PTSD status did little to predict risk of alcohol use and abuse, after the influence of other variables was controlled.

Being older, Caucasian, male, and from higher SES levels also increased risk of marijuana abuse or dependence, when all other variables were considered. Familial drug use was strongly associated with use and abuse of marijuana in adolescents, but familial alcohol abuse was not consistently related to problematic marijuana use. Physical and sexual abuse more than doubled risk of problem use of marijuana, independent of effects of familial substance use. Once again, witnessed violence was also strongly associated with problem marijuana use. Independent effects of current PTSD status on past-year marijuana problem use were also notable, with PTSD-positive individuals at twice the risk of abuse.

When all variables were considered in the final model, race was strongly related to past year hard drug use and problem use, with Caucasians at 2-3 times the risk. Age was also positively associated with risk. Familial drug and alcohol use increased risk of hard drug problem use by a factor of 4, whereas sexual and physical abuse were associated with a large increase in risk, over and above effects of familial substance use. Once again, witnessed violence led to increased rates of recent use and problem use, while PTSD status had little impact independent that of other variables. The consistent relationship between PTSD diagnostic status and marijuana problem use was not observed for hard drugs (or alcohol).

Univariate and multivariate analyses in which substance use was the predicted variable were conducted, this time substituting current delinquency as the predicted variable. The same set of variables used to predict substance abuse was employed to predict delinquency (with the addition of substance abuse as a predictor variable, of course). Multivariate analyses are primarily considered here.

For male adolescents, age, race, and income were not associated with changes in final model risk of delinquency. However, familial drug or alcohol problems doubled risk of delinquency, over and above effects produced by race and income. The final model further revealed that physical assault, but not sexual assault increased risk that males would engage in delinquent acts. This may be an artifact of the small number of male subjects reporting sexual abuse, however. Witnessed violence and problem substance use also greatly increased risk of delinquency, but this was expected given the definitional overlap between juvenile delinquency behavior and these predictors. Finally, being diagnosed with PTSD slightly increased the likelihood that male adolescents would report engaging in delinquent acts.

Considering again the final model, for female respondents, age was inversely related to risk of delinquency. Familial drug and alcohol abuse doubled risk of delinquency, as did physical, but not sexual assault. Witnessed violence more than tripled risk of delinquency, and problem substance use increased risk by a factor of 6. For girls, PTSD also increased risk of engaging in illegal activity.

From these analyses, it is clear that the pathways to delinquency, at least based on the factors assessed here, may vary by gender and ethnicity. Specifically, African-American males seem to have less in common with White males or African-American females than might be expected. Several of the significant predictors for these other groups (e.g., family history of problems with substance use variables and history of physical assault or physical abuse) were not relevant for African-American males. Also, a history of PTSD was important for African-American males, but not for the other group. Therefore, these adolescents may have a unique set of circumstances that lead to delinquency.

Overall, logistic regression analyses indicated that the key variables in predicting delinquency status were substance use, victimization history, particularly physical assault and familial substance use. PTSD status also added somewhat to prediction of delinquency status. Perhaps the most important result of these analyses was the finding that any variable in isolation, including substance use, had limited impact on delinquency outcome. Indeed, substance use, history of victimization, or family history of substance abuse were all associated with approximately the same levels of delinquency. Combinations of variables, however, yielded a very different picture. Fully 78% of adolescents who were substance abusers, had been victimized, and had family members that used substances engaged in past-year delinquency. However, adolescents with all three risk factors represented only 15% of the male and 27% of the female delinquent population. Attention was therefore appropriately directed to pairs of variables. Levels of delinquency in children with both victimization and substance use histories or both family substance use and personal substance use histories, or both victimization and family substance use histories were tremendously elevated, and, of equal importance, each of

these combinations of variables accounted for 30% to 50% of the delinquent population. Thus, as outlined above, (a) testing positive for only two of the risk factors greatly increases the likelihood that a child will engage in delinquent behavior, and (b) this is very important because a large proportion of delinquency in this country appears to be etiologically related to these variables.

Assessment and treatment implications are clear: measurement of substance use in delinquents must be complemented by measurement of victimization history and familial substance abuse. Adolescent substance use alone does not determine delinquency, but is itself determined, in large part, by variables that also contribute to prediction of delinquency (i.e., victimization and familial substance use). Treatment for delinquency should follow assessment findings, and strategies to reduce substance use should be combined with techniques to diminish negative effects of victimization and familial substance use in children with these histories. In so doing, both adolescent substance use and delinquency will be more effectively reduced.

Summary

Overall, observed patterns of association between relevant variables and substance use and problem use were remarkably consistent across drug types, racial backgrounds, incomes, and gender. Risk of use and abuse of all substances in general, but hard drugs in particular, was greatly increased in adolescents who had suffered sexual assault, physical assault and abuse, or who had witnessed violence. Familial substance use was also a source of consistent and significant risk. Notably, effects of victimization were independent those of other variables, further supporting the role of sexual or physical assault of adolescents in facilitating development of substance use and abuse. Moreover, the temporally linear contribution of assault to substance use behavior is indicated by the finding that victimization preceded substance use in most cases. Importantly, the overt symptom constellation that comprises post-traumatic stress disorder did not consistently elevate risk of alcohol or hard problem use, when demographic and victimization variables were controlled. Thus, even though victimized adolescents may not display prototypical adult post-trauma symptoms, they appear to be at high risk of suffering significant negative effects of trauma, particularly in the form of substance use disorders. This problem is chronological in nature, and ironically, sets the stage for future victimization (Kilpatrick, Acierno, Resnick, Saunders, & Best, 1997).

Three factors were consistently associated with increased delinquency: 1) victimization, 2) substance use, and 3) familial substance use. Substance abuse alone had limited impact on delinquency status, relative to the combined presence of victimization and family substance abuse histories. Presence of all three variables tremendously increased risk of delinquency, but most delinquents (over 85%) did not test positive for all three variables. However, testing positive for any two variables also led to greatly elevated rates of delinquency, and more importantly, described over 50% of the juvenile population. Thus, relevance of victimization history and familial substance use, in addition to adolescent substance abuse to delinquency status, was demonstrated.

VII. RECOMMENDATIONS FOR RESEARCH, POLICY, AND PRACTICE

A. Recommendations for Research:

As is the case with most research, results of the NSA raise as many questions as they answer. Rather than identifying a long list of recommendations for addiction research that could be generated by a review of the NSA findings, we present five recommendations that represent the highest priorities for future research.

Recommendation One: Longitudinal research is needed to clarify the temporal sequence of victimization, PTSD, substance use/abuse/dependence, and delinquent behavior among adolescents. This is particularly important given that rates of violent assault, witnessing violence, alcohol and drug use, and delinquent behavior between the ages of 12 and 17. Such research should examine the temporal sequence of problem development as well as risk and protective factors that are related to victimization, alcohol and drug use, PTSD, and delinquent behavior.

Recommendation Two: The NSA demonstrated the feasibility of obtaining information from adolescents about victimization experiences. Researchers should be encouraged to include measures for screening for history of violent assault and witnessing violence in studies of adolescent alcohol and drug use and delinquency.

Recommendation Three: NSA findings indicate that PTSD appears to be a mediating factor in the relationship between victimization and substance use/abuse/dependence problems between victimization and delinquent behavior. Therefore, research should be conducted examining the efficacy of preventative mental health treatments for PTSD on the subsequent development of substance use/abuse/dependence and delinquent behavior problems.

Recommendation Four: NSA findings indicate that the bulk of violent assaults are perpetrated by someone the victim knows well rather than by a stranger. Future research should obtain more information about the circumstances and behavioral sequences that precede and follow such assaults. This might provide valuable data that would prove useful in the design of violence prevention programs.

Recommendation Five: A longitudinal follow-up study should be done with the NSA sample.

Recommendation Six: Research should examine specific and separate analytical models for gender and racial/ethnic groups. Results of the NSA indicated that not only were there significant differences between gender and racial/ethnic groups on the prevalence of many major variables, but relationships between these variables varied between these subgroups. For example, PTSD appeared to be a more important predictor of delinquency for African-American males than for other gender-racial/ethnic subgroups. Therefore, it is recommended that predictive models and other analyses be conducted by and within gender and racial/ethnic groups to further expose and understand these subgroup differences.

Recommendation Seven: The roles of specific types of victimization and particular characteristics of victimizations should be evaluated in the development of substance use problems and delinquency, especially with gender and racial ethnic subgroups. Some results suggested that some types of victimization are more important for some subgroups for predicting these problems. Further refinement and testing of these hypotheses are needed.

Recommendation Eight: The recommendations above suggest that larger sample sizes and/or purposive sampling methods are needed to conduct more specific and refined research among important subgroups. In order to achieve the cell sample sizes necessary to understand the roles of specific types of victimizations with certain characteristics among gender and racial ethnic subgroups, larger initial sample sizes will be necessary. Alternatively, purposive sampling methods could be used to increase cell sample sizes while maintaining the level generalizability and external validity necessary for meaningful results.

Recommendation Nine: Research should be conducted to better understand the factors that contribute to the dramatic under-reporting of crimes against children.

While some research exists in this area, most reasons offered for under-reporting is simply conjecture. Intervention (and secondary and tertiary prevention) cannot occur without identification.

B. Recommendations for Policy:

Recommendation One: The Bureau of Justice Statistics should consider making several changes in the National Crime Victimization Survey based on the NSA findings. Adolescents should be asked about sexual and physical assaults using more explicit screening questions along the lines of those demonstrated to be feasible in the NSA., Likewise, the NCVS should be revised to include brief measures of crime-related mental health problems

Recommendation Two: The level of peer violence documented by the NSA suggests that much of the violence adolescents experience is perpetrated by other adolescents, many of whom would be processed by the juvenile justice system rather than by the criminal justice system. Thus, the juvenile justice system must be upgraded to insure that adolescent victims of violence perpetrated by juveniles receive comparable victim assistance as victims in the criminal justice system.

Recommendation Three: The extent to which violent assault go unreported to criminal justice system authorities suggests that it is important to identify barriers to reporting as well as ways to increase reporting to authorities.

Recommendation Four: The NSA found that many violence victims had curved PTSD, and substance use/abuse/dependence problems. This suggests that these problems are persistent among victims who do not get effective mental health treatment. Therefore, mechanisms should be developed to insure that funding is available to provide mental health counseling to adolescent victims who need it irrespective of their ability to pay or whether they qualify for crime victim compensation.

□ Recommendation Five: A greater emphasis should be placed on youth victimization in juvenile justice policy and programs. Results of the NSA strongly indicate that victimization and its mental health correlates play an important role in the development of substance use and delinquency behavior among many adolescents. However, victimization receives relatively little attention in juvenile justice public policy compared to offending behavior. This small level of policy attention is shortsighted and neglects a key risk factor for these problems. Therefore, a reorientation in policy attention is required. The role of child victimization and its effects should be considered in all juvenile justice policy and program initiatives.

□ Recommendation Six: Policies should promote the prevention of child victimization as part of a comprehensive plan for preventing youth substance use and delinquency. Effective and efficient prevention begins as early as possible in the risk factor chain. Results of this study suggest that victimization and its effects are strong and primary correlates with youth substance abuse and delinquency. Therefore, prevention of these early primary experiences will contribute to preventing these secondary problems.

□ Recommendation Seven: Policies should encourage early identification of and intervention with victimized children (secondary and tertiary prevention). All child victimizations cannot be prevented. However, if more can be recognized and effective intervention provided to child victims, it is likely that at least some of the long-term negative effects leading to substance use and delinquency can be mitigated. Therefore, policies should encourage proactive rather than reactive approaches to identifying victimized youth, and should promote providing effective and rapid intervention for victimization-related problems that are related to the development of substance use and delinquency.

C. Recommendations for Practice

□ Recommendation One: Mental health professionals who work with children and adolescents should be informed about the high rates of victimization that occur among children and adolescents and about the extent to which victimization serves as a risk factor for PTSD, substance use/abuse/dependence, and delinquency.

□ Recommendation Two: Mental health professional should be encouraged to screen for victimization experiences among child and adolescent clients. Substance abuse treatment programs for adolescents should do likewise.

□ Recommendation Three: Victim assistance professionals in the criminal and juvenile justice systems should establish relationships with mental health professionals who are knowledgeable about crime victims' mental health issues.

□ Recommendation Four: Mental health programs dealing with child victims should incorporate substance abuse and delinquency prevention components. While mental health programs designed to reduce common psychological problems associated with child victimization are common, few include specific interventions for reducing substance use

onset, substance abuse, or conduct and delinquency problems. Given the findings of the NSA, mental health programs should incorporate these prevention components as a regular part of their victimization treatment protocols.

□ Recommendation Five: Law enforcement should develop programs for dramatically increasing the number of crimes against children that are reported, identified and investigated by them. For example, some departments have developed specialized Crimes Against Children unit in order to develop and focus specialized expertise on this area of crime. While the effectiveness of such initiatives is unknown, their purpose is important. Creative law enforcement attention is needed to stop the current crime wave against children. By limiting the number of crimes committed AGAINST children, the number of crimes committed BY children likely will decrease. Unfortunately, unless the crimes are identified, they cannot be investigated. Proactive, creative community programs are needed to encourage children and others to report crimes to law enforcement. It is likely that if the reporting rate of crimes against children remains low, the crime wave against children will continue, and more children will go on to commit crimes.

□ Recommendation Six: Prosecutors should develop specific programs and expertise to increase the proportion of crimes against children that are successfully prosecuted. Many offices have tried various programs. The effectiveness of these programs is unknown for the most part. However, it is likely that if prosecutors focus more attention on crimes with child victims, more will be successfully prosecuted. More successful prosecution likely means fewer crimes against children.

□ Recommendation Seven: Victim/witness programs should develop specific programs and expertise for dealing with child and adolescent victims and witnesses in order to encourage increased reporting and cooperation with the criminal justice system.

VIII. REFERENCES

- Agarwal, S.N. (1983). *Sexual assault among adolescents*. Lexington, MA: Lexington Books.
- Bastian, L.D., & Taylor, B.M. (1991). *School crime: A National Crime Victimization Survey Report* (NCJ-131645). Washington, DC: U. S. Department of Justice.
- Beitchman, J.H., Zucker, K.J., Hood, J.E., DaCosta, G.A., Akman, D., & Cassavia, E. (1992). A review of the long-term effects of child sexual abuse. *Child Abuse & Neglect*, 16, 101-113.
- Boyle, J.M., & Kilpatrick, D.G. (May, 1993). *Non-participation bias in telephone survey of children*. Paper presented at the 1993 Annual Conference of American Association for Public Opinion Research, St. Charles, IL.
- Bradburn, N.M. (1984). *Discussion: Telephone survey methodology*. Proceedings of the 4th Conference on Health Survey Research Methods, Washington, DC [DHHS Publication No. (PHS) 84-3346]. U.S. Department of Health and Human Services.
- Brady, K.T., Killeen, T., Saladin, M., Dansky, B.S., & Becker, S. (1994). Cocaine, substance abuse and PTSD: Characteristics of women in treatment. *American Journal of Addictions*, 3, 160-164.
- Brook, J.S., Nomura, C., & Cohen, P. (1989). A network of influences on adolescent drug involvement: Neighborhood, school, peer, and family. *Genetic, Social and General Psychology Monographs*, 115, 125-145.
- Brook, J.S., Whiteman, M.M., & Finch, (1992). Childhood aggression, adolescent delinquency, and drug use: A longitudinal study. *The Journal of Genetic Psychology*, 155, 369-383.
- Browne, A., & Finkelhor, D. (1986). Impact of child sexual abuse: A review of the research. *Psychological Bulletin*, 99, 66-77.
- Burnam, M.A., Stein, J.A., Golding, J.M., Siegel, J.M., Sorenson, S.B., Forsythe, A.B., & Tefles, C.A. (1988). Sexual assault and mental disorders in a community population. *Journal of Consulting and Clinical Psychology*, 6, 843-850.
- Catlin, G., & Murray, S. (1979). *Report on Canadian victimization survey methodology, pretests*. Ottawa: Statistics Canada.
- Centers for Disease Control (1990). Behavioral risk factor surveillance (1988). *Morbidity and mortality weekly report*, 39 (Supplement No. SS-2), 1-22.

Cottler, L.B., Compton, W.M., Mager, D., Spitznagel, E.L., and Janca, A. (1992). Posttraumatic stress disorder among substance users from the general population. *American Journal of Psychiatry*, 149, 644-647.

Cottler, L.B., Compton, L.B., Mager, D., Spitznagel, E.L., & Janca, A. (1992). Posttraumatic stress disorder among substance users from the general population. *Journal of Consulting and Clinical Psychology*, 56, 843-850.

Dembo, R., Dertke, M., LaVoie, L., Borders, S., Washburn, M., & Schmeidler, J. (1987). Physical abuse, sexual victimization and illicit drug use: A structural analysis among high risk adolescents. *Journal of Adolescence*, 10, 13-33.

Dembo, R., Williams, L., La Voie, L., Berry, E., Getreu, A., Wish, E.D., Schmeidler, J., & Washburn, M. (1989). Physical abuse, sexual victimization, and illicit drug use: Replication of a structural analysis among a new sample of high-risk youths. *Violence and Victims*, 4, 121-138.

Dembo, R., Williams, L., Schmeidler, J., Berry, E., Wothke, W., Getreu, A., Wish, E.D., & Christensen, C. (1992). A structural model examining the relationship between physical child abuse, sexual victimization, and marijuana/hashish use in delinquent youth: A longitudinal study. *Violence and Victims*, 7, 41-62.

Dembo, R., Williams, L., Wish, E.D., Dertke, M., Berry, E., Getreu, A., Washburn, M., & Schmeidler, J. (1988b). The relationship between physical and sexual abuse and illicit drug use: A replication among a new sample of youths entering a juvenile detention center. *International Journal of the Addictions*, 22, 1101-1123.

Elliott, D. M. (1992). *Traumatic Events Survey*. Unpublished psychological test, Harbor-UCLA Medical Center, Los Angeles.

Elliott, D., 1994. personal communication.

Elliott, D., Huizinga, D., & Ageton, S. S. (1985). *Explaining Delinquency and Drug Abuse*. Sage Publications.

Elliott, D.S., Huizinga, D., & Menard, S. (1988). *Multiple problem youth, delinquency, substance use, and mental health problems*. New York: Springer-Verlag

Elliott, D., Huizinga, D., & Menard, S. (1989). *Multiple problem youth: Delinquency, substance use, and mental health problems*. New York: Springer-Verlag.

Finklehor, D. & Dziuba-Leatherman, J. (1994). Victimization of Children. *American Psychologist*, 49(3), 173-183.

Finkelhor, D., Hotelling, G., Lewis, I.A., & Smith, C. (1990). Sexual abuse in a national survey of adult men and women: Prevalence, characteristics, and risk factors. *Child Abuse & Neglect*, 14, 29-38.

Gelles, R.J., & Straus, M.A. (1987). Is violence toward children increasing?: A comparison of 1975 and 1985 national survey rates. *Journal of Interpersonal Violence*, 2(2), 212-222.

Gelles, R.J., & Straus, M.A. (1990). The medical and psychological costs of family violence. In M.A. Straus & R.J. Gelles (Eds.), *Physical violence in American families: Risk factors and adaptations to violence in 8,145 families*. New Brunswick, NJ: Transaction, pp. 425-430.

George, L.K., & Winfield-Laird, I. (1986). *Sexual assault: Prevalence and mental health consequences*. Final report submitted to NIMH.

Hanson, R.F., Kilpatrick, D.G., Freedy, J.R., Saunders, B.E. (1995). Los Angeles County following the civil disturbances: Degree of exposure and impact on mental health. *Journal of Consulting and Clinical Psychology*, 63(6), 987-996.

Helzer, J.E., Robins, L.N., & McEvoy, L. (1987). Post-traumatic stress disorder in the general population. *New England Journal of Medicine*, 317, 1630-1634.

Huizinga, D., Loeber, R., & Thornberry, T.P. (1995). *Urban Delinquency and Substance Abuse: Initial Findings* (OJJDP-86-JN-CX-006,007,008). Washington, DC: Office of Justice Programs.

Jaffe, P.G., Wolfe, D.A., & Wilson, S.K. (1990). *Children of Battered Women*. Newbury Park: Sage.

Johnston, L.D., O'Malley, P.M., & Bachman, J.G. (1995). *National survey results on drug use from the Monitoring the Future study, 1975-1994. Volume I: Secondary school students* (NIH Pub. No. 95-4026). *Volume II: College students and young adults* (1996). (NIH Pub. No. 96-4027). Rockville, MD: National Institute on Drug Abuse. 327 pp. and 189 pp., respectively.

Kendall-Tackett, K.A., Williams, L.M., & Finkelhor, D. (1993). Impact of sexual abuse on children: A review and synthesis of recent empirical studies. *Psychology Bulletin*, 113, 164-180.

Kessler, R.C., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C.B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*, 52, 1048-1060.

Kilpatrick, D.G. (1983). Rape victims: Detection, assessment, and treatment. *The Clinical Psychologist*, 36(4), 92-95.

Kilpatrick, D.G. (1990, August). Violence as a precursor of women's substance abuse: The rest of the drugs-violence story. Presentation at the 98th annual convention of the American Psychological Association, Boston, MA.

Kilpatrick, D.G. (March, 1992). *More than you ever wanted to know about crime victims and their mental health problems*. Invited address at the 38th Annual Meeting of the Southeastern Psychological Association, Knoxville, TN.

Kilpatrick, D.G., Edmunds, C.N., & Seymour, A.K. (1992). *Rape in America: A report to the nation*. Arlington, VA: National Victim Center.

Kilpatrick, D.G. & Resnick, H.S. (March, 1994). *Rape, other violence against women, and post-traumatic stress disorder: Critical issues in assessing the adversity-stress-psychopathology relationship*. 84th annual meeting of the American Psychopathological Association, New York.

Kilpatrick, D.G. & Resnick, H.S. (Sept., 1994). *Victimization and Post Traumatic Stress Disorder*. Presented at a conference sponsored by NIDA's Office of the Director entitled "Drug Addiction Research and Health of Women", Washington, DC.

Kilpatrick, D.G., Resnick, H.S., Freedy, J.R., Pelcovitz, D., Resick, P.A., Roth, S., & van der Kolk, B. (1994). The posttraumatic stress disorder field trial: Emphasis on Criterion A and overall PTSD diagnosis. *DSM-IV Sourcebook*. Washington, DC: American Psychiatric Press.

Kilpatrick, D.G., Resnick, H.S., Saunders, B.E. & Best, C.L. (in press). Victimization, PTSD, and substance use/abuse among women. *Drug Addiction Research and Health of Women*.

Kilpatrick, D.G., Resnick, H.S., Saunders, B.E., Best, C.L., & Epstein, J. (June, 1994). *Violent assault and alcohol dependence among women: Results of a longitudinal study*. Poster presented at the annual meeting of the Research Society on Alcoholism, Maui, Hawaii.

Koss, M.P. (1993). Detecting the scope of rape: A review of prevalence research methods. *Journal of Interpersonal Violence*, 8(2), pp. 198-222.

Kulka, R.A., Schlenger, W.E., Fairbank, J.A., Hough, R.L., Jordan, B.K., Marinar, C.R., & Weiss, J.A. (1990). *Trauma and the Vietnam war generation: Report of findings from the National Vietnam Veterans readjustment study*. New York: Brunner/Mazel.

Ladwig G.B. & Anderson, M.D. (1989). Substance abuse in women: Relationship between chemical dependency of women and past reports of physical/or sexual abuse. *International Journal of the Addictions* 24(8), pp. 739-754.

Franktree, C.B., Briere, J., & Zaidi, L. (1991). Incidence and impact of sexual abuse in a community sample: The role of direct inquiry. *Child Abuse and Neglect*, 15, 447-458.

Lewis, D.O., Shanok, S.S., Pincus, J.H., & Glaser, G.H. (1979). Victimized juvenile delinquents: Psychiatric, neurological, psychological, and abuse factors. *Journal of the American Academy of Child and Adolescent Psychiatry*, 18, 307-319.

Lipovsky, J.A., Saunders, B.E., & Murphy, S.M. (1989). Depression, anxiety, and behavior problems among victims of father-child sexual assault and nonabused siblings. *Journal of Interpersonal Violence*, 4, 452-468.

McCurdy, K., & Daro, D. (1993). *Current trends in child abuse reporting and treatment: the results of the 1992 Annual Fifty State Survey*. Chicago, IL: National Committee for Prevention of Child Abuse.

Miller, B.A., Downs, W.R., Gondoli, D.M., and Keil, A. (1987). The role of childhood sexual abuse in the development of alcoholism in women. *Violence and Victims*, 2(3), 157-172.

Miller, B.A., Wieczorek, W.F., Downs, W.R. (June, 1994). *Victimization experiences and problem drinking among five samples of women*. Presented at the Research Society on Alcoholism, Maui, HI.

Murphy, S.M., Amick-McMullan, A., Kilpatrick, D.G., Haskett, M.E., Veronen, L., Best, C.L., & Saunders, B.E. (1988). Rape victim's self-esteem: A longitudinal analysis. *Journal of Interpersonal Violence*, 1(4), 355-370.

National Center on Child Abuse and Neglect, 1994 National Household Survey of Drug Abuse (1995 Paulsen, A.S., Crowe, R.R., Noyes, R., & Pfohl, B. (1988). Reliability of the telephone interview in diagnosing anxiety disorders. *Archives of General Psychiatry*, 45, 62-63.

Paulsen, A.S., Crowe, R.R., Noyes, R., & Pfohl, B. (1988). Reliability of the telephone interview in diagnosing anxiety disorders. *Archives of General Psychiatry*, 45, 62-63.

Projections of the Population of States by Age, Sex and Race: 1988 to 2010. *Current Population Reports*, P-25, No. 1017, 1988.

Resnick, H.S., Kilpatrick, D.G., Dansky, B.S., Saunders, B.E., & Best, C.L. (1993). Prevalence of civilian trauma and PTSD in a representative national sample of women. *Journal of Consulting & Clinical Psychology* 61(6).

Saunders, B.E., Kilpatrick, D.G., Lipovsky, J.A., Resnick, H.S., Best, C.L., & Sturgis, E.T. (1991, March). *Prevalence, case characteristics, and long-term psychological effects of child sexual assault: A national survey*. Presentation at the annual meeting of the American Orthopsychiatric Association, Toronto, Ontario, Canada.

1987) Child sexual assault as a risk factor for mental disorders among women: A community study. *Journal of Interpersonal Violence*, 2(2), 184-204.

Sorenson, S.B., Stein, J.A., Burnam, M.A., Golding, J.M., Brown, & Siegel, J.M. (1987). Prevalence of adult sexual assault: The Los Angeles Epidemiologic Catchment Area Study. *American Journal of Epidemiology*, 126, 1145-1164.

Sorenson, S.B., Stein, J.A., Siegel, J.M., Golding, J.M. & Burnam, M.A. (1987). The prevalence of adult sexual assault: The Los Angeles epidemiologic catchment area project. *American Journal of Epidemiology*, 126, 1154-1164.

Stein, J.A., Golding, J.M., Siegel, J.M., Burnam, M.A., & Sorenson, S.B. (1988). Long-term psychological sequelae of child sexual abuse: The Los Angeles Epidemiologic Catchment area study. In G. E. Wyatt & G.J. Powell (Eds.), *Lasting effects of child sexual abuse*. Newbury Park: Sage Publications, pp. 135-154.

Straus, M.A. (1984). Family violence and non-family crime and violence. In A.J. Lincoln & M.A. Straus (Eds.), *Crime and the Family*, Springfield, IL: Thomas.

Thornberry, T. (1994). Violent families and youth violence. *OJJDP Fact Sheet*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention.

Von, J.M., Kilpatrick, D.G., Burgess, A.W., & Hartman, C.R. (1991). Rape and sexual assault. In M.L. Rosenberg & M.A. Fenley (Eds.), *Violence in America: A public health approach* (pp. 95-122). New York: Oxford University Press.

Weeks, M.F., Kulka, R.A., Lessler, J.T., & Whitmore, R.W. (1983). Personal versus telephone surveys for collecting household health data at the local level. *American Journal of Public Health*, 73, 1389-1394.

Whitaker, C.J., & Bastian, L.D. (1991). *Teenage Victims: A National Crime Survey Report*. Washington, D.C.: United States Department of Justice

Widom, C.S. (1989). The cycle of violence. *Science*, 244, 160-166.

Widom, C.S. (1992). The cycle of violence. *National Institute of Justice: Research in Brief*. Washington, D.C.: U.S. Department of Justice.

Table 1. Demographic Characteristics of Parent/Guardian Sample

	n	Percent
Gender		
Female	2887	71.8
Male	1136	28.2
Relationship to the Adolescent		
Biological Parent	3622	90.0
Step-parent	201	5.0
Adoptive Parent	60	1.5
Grandparent	54	1.3
Another Relative	26	0.6
Guardian	42	1.0
Something else	17	0.4
Refused	1	0.0
Marital Status		
Married	3096	77.0
Living as couple	143	3.5
Separated	149	3.7
Divorced	401	10.0
Widowed	61	1.5
Single/never married	166	4.1
Refused	7	0.2

Table 1 (con't)

Demographic Characteristics of Parent/Guardian Sample

Occupational Status		
Employed full-time	3178	79.0
Employed part-time	284	7.1
In the military	22	0.5
Unemployed & looking for work	122	3.0
Retired	68	1.7
Student	33	0.8
Disabled or too ill to work	70	1.7
Other	65	1.6
Refused	1	0.0
Total Household Income		
More than \$50,000	1362	33.9
\$30,001 to \$50,000	1168	29.0
\$20,001 to \$30,000	600	14.9
\$10,001 to \$20,000	368	9.1
\$5,000 or less	114	2.8
Not sure/refused	232	5.8
Highest Educational Achievement		
Graduate degree	426	10.6
Some Graduate school	131	3.3
Four year college graduate	658	16.3
Some college	1100	27.4

Table 1 (cont.)

Demographic Characteristics of Parent/Guardian Sample

Highest Educational Achievement		
High School graduate	1288	32.0
Some high school	316	7.8
Eighth grade	54	1.3
First through seventh grade	39	1.0
No formal schooling	1	0.0
Not sure/Refused	10	0.2
Racial/Ethnic Identity		
Caucasian/Non-Hispanic	3055	75.9
African American/Non-Hispanic	592	14.7
Hispanic	270	6.7
Native American	31	0.8
Asian	30	0.8
Something else	31	0.8
Not sure/refused	14	0.3
Community of Residence		
Large city	639	15.9
Suburb of large city	764	19.0
Large town	735	18.3
Small town	1095	27.2
Rural area	783	19.5
Not sure/refused	7	0.2

NOTE: N = 4,023

Table 2. Demographic Characteristics of Adolescent Sample

	N	Percent
Gender		
Female	1958	48.7
Male	2065	51.3
Racial/Ethnic Identity		
Caucasian/Non-Hispanic	2825	70.2
African American/Non-Hispanic	590	14.7
Hispanic	314	7.8
Native American	139	3.5
Asian	46	1.1
Other	81	2.0
Not Sure/Refused	28	0.7
Age		
12	682	17.0
13	685	17.0
14	673	16.7
15	682	16.9
16	652	16.2
17	641	15.9
Not sure/Refused	8	0.2
Year in School		
Fifth	44	1.1
Sixth	284	7.1
Seventh	711	17.7

Table 2 (cont.)

Demographic Characteristics of Adolescent Sample

Year in School

Eighth	659	16.4
Ninth	745	18.5
Tenth	643	16.0
Eleventh	561	13.9
Twelfth	330	8.2
Don't Attend	43	1.1
Not Sure/Refused	3	0.1

Note: $N = 4,023$

Table 3. Lifetime Prevalence of Types of Sexual Contact by Gender

Act	GENDER		SIG
	Male	Female	Level
Penile penetration of child	0.5%	3.3%	.0001
Finger/Object penetration of child	0.6%	2.7%	.0001
Others' mouth on child's sexual parts	1.0%	1.3%	NS
Touching of child's sexual parts	2.8%	9.9%	.0001
Child forced to touch others' sexual parts	0.7%	3.5%	.0001
Unwanted penetration of others by child (asked only of males)	0.8%	n/a	n/a
Any Sexual Assault	3.4%	13.0%	.0001

Table 4. Lifetime Prevalence of Physical Assault by Gender

Event	GENDER		SIG
	Male	Female	Level
Attacked with weapon	6.0%	3.4%	.002
Attacked with intent to kill/injure	8.5%	6.7%	NS
Threatened with gun or knife	7.9%	4.3%	.0001
Beaten w/object, hurt badly	5.9%	3.5%	.004
Beaten w/fists, hurt badly	7.4%	5.1%	.01
Any physical assault	21.3%	13.4%	.0001

Table 5. Lifetime Prevalence of Physically Abusive Punishment by Gender

Event	GENDER		SIG
	Male	Female	Level
Spanked so hard you had to see a doctor	0.2%	0.7%	.05
Spanked to hard you got bad marks, bruises, cuts, or welts	8.2%	9.9%	ns
Punished by burning, cutting or tying you up	0.6%	0.4%	ns
Any physically abusive punishment	8.5%	10.2%	ns

Table 6. Lifetime Prevalence of Witnessing Violence by Gender

Event	GENDER		SIG
	Male	Female	Level
Seen someone shot with a gun	5.8%	4.1%	.01
Seen someone stabbed/cut	12.1%	9.0%	.01
Seen someone sexually assaulted	1.8%	3.8%	.001
Seen someone mugged/robbed	14.8%	7.8%	.0001
Seen someone threatened with a weapon	38.0%	28.6%	.0001
Any witnessing violence	43.6%	35.0%	.0001

Table 7. Initial and Final Model Odds of Lifetime PTSD as a Function of Demographic Variables, Family Substance Use, and Sexual Assault, Physical Assault, and Witnessed Violence: Hierarchical Logistic Regression Analyses

<u>Step</u>	<u>Variable</u>	<u>Beta</u>	<u>SE</u>	<u>Wald</u>	<u>Step OR</u>	<u>Final OR</u>	<u>Final p</u>
1	Age	.25	.04	45.5	1.28	1.15	.001
	Gender (female)	.52	.12	18.1	1.69	1.82	.000
	Race (Nonwhite)	.29	.14	4.5	1.34	ns	.718
	Income	-.06	.03	3.1	ns	ns	.559
2	Family Alcohol Problem	1.04	.14	52.3	2.82	1.60	.003
	Family Drug Problem	.86	.16	28.1	2.37	1.81	.001
3	Number of Sexual Assaults	.53	.11	22.9	1.69	1.69	.000
	Number of Physical Assaults	.56	.09	36.6	1.75	1.75	.000
	Number Witnessed Violence	.65	.09	52.1	1.92	1.92	.000

Note: Beta, SE, Wald and Final p are given for the final model.

Table 8: Univariate Odds of Past-Year Alcohol Abuse/Dependence, Marijuana Abuse/Dependence, and Hard Drug Abuse/Dependence as a Function of Demographic Variables, Familial Problem Alcohol Use, Familial Problem Drug Use, Physical Abuse/Assault, Sexual Assault, Witnessed Violence, and Current PTSD: Chi Square Analyses

<u>Analysis A. Odds of Alcohol Abuse/Dependence</u>			
<u>Variable</u>	<u>Odds Ratio</u>	<u>χ^2/Wald</u>	<u>p</u>
Age ^a	2.03	110.4	.000
Gender ^b	ns	4.03	.045
Race ^b	ns	2.16	.117
Income	1.14	9.41	.002
Familial Problem Alcohol	3.67	60.0	.000
Familial Drug Use	1.93	7.70	.006
Physical Abuse/Assault	4.00	79.7	.000
Sexual Assault	4.65	75.0	.000
Witnessed Violence	4.89	84.9	.000
Current PTSD	4.00	38.8	.000
<u>Analysis B. Odds of Marijuana Abuse/Dependence</u>			
Age ^a	1.65	71.4	.000
Gender ^b	ns	4.71	.029
Race ^b	ns	1.20	.273
Income	ns	3.51	.060
Familial Problem Alcohol	3.33	46.2	.000
Familial Drug Use	4.12	59.2	.000
Physical Abuse/Assault	4.92	102.2	.000
Sexual Assault	3.84	48.2	.000
Witnessed Violence	8.58	124.7	.000
Current PTSD	6.20	88.1	.000

Note: Bonferroni correction requires $p < .01$ to maintain setwise alpha at .096.

Table 8 continued.

Analysis C. Odds of Hard Drug Abuse/Dependence

<u>Variable</u>	<u>Odds Ratio</u>	<u>χ^2 / Wald</u>	<u>p</u>
Age ^a	1.81	22.3	.000
Gender ^b	ns	0.00	1.00
Race ^b	ns	0.67	.413
Income	ns	2.74	.098
Familial Problem Alcohol	7.87	50.0	.000
Familial Drug Use	7.93	48.6	.000
Physical Abuse/Assault	12.44	62.8	.000
Sexual Assault	8.73	54.8	.000
Witnessed Violence	13.38	37.3	.000
Current PTSD	8.80	45.7	.000

Note: Bonferroni correction requires $p < .01$ to maintain setwise alpha at .096.

^a Continuous variables were analyzed through single-predictor logistic regression and the Wald statistic is reported. ^b Odds ratios greater than 1 for these variables indicate that being male or being Caucasian was associated with increased risk.

Table 9: Initial and Final Model Odds of Past-Year Alcohol Abuse/Dependence, Marijuana Abuse/Dependence, and Hard Drug Abuse/Dependence as a Function of Demographic Variables, Familial Problem Alcohol Use, Familial Problem Drug Use, Physical Abuse/Assault, Sexual Assault, Witnessed Violence, and Current PTSD: Hierarchical Logistic Regression Analyses

<u>Regression A. Odds of Alcohol Abuse/Dependence</u>							
<u>Step</u>	<u>Variable</u>	<u>Beta</u>	<u>SE</u>	<u>Wald</u>	<u>Step OR</u>	<u>Final OR</u>	<u>Final p</u>
1	Age	.66	.072	83.3	2.06	1.93	.000
	Gender ^a	-.50	.195	6.48	1.42	1.64	.011
0	Race ^a	-.63	.237	7.04	ns	1.88	.008
	Income	.18	.049	14.1	1.12	1.20	.000
2	Familial Problem Alc.	.86	.208	16.8	3.51	2.35	.000
	Familial Drug Use	-.10	.269	.137	ns	ns	.712
3	Physical Abuse/Asslt.	.59	.204	8.26	2.42	1.80	.004
	Sexual Assault	.94	.246	14.6	2.95	2.56	.001
4	Witnessed Violence	.96	.215	19.9	2.62	2.60	.000
5	PTSD	.19	.288	.436	ns	ns	.509
<u>Regression B. Odds of Marijuana Abuse/Dependence</u>							
1	Age	.42	.065	41.6	1.66	1.52	.000
	Gender ^a	-.52	.200	6.64	1.49	1.67	.010
	Race ^a	-.56	.229	5.89	ns	1.74	.015
	Income	.13	.049	7.33	ns	1.14	.007
2	Familial Problem Alc.	.30	.217	1.96	2.17	ns	.162
	Familial Drug Use	.80	.231	12.04	3.38	2.23	.001
3	Physical Abuse/Asslt.	.51	.205	6.08	2.71	1.66	.013
	Sexual Assault	.56	.255	4.90	2.46	1.76	.027
4	Witnessed Violence	1.51	.247	37.6	4.73	4.56	.000
5	PTSD	.87	.255	11.7	2.39	2.39	.001

Table 9 continued.

<u>Regression C. Odds of Hard Drug Abuse/Dependence</u>							
<u>Step</u>	<u>Variable</u>	<u>Beta</u>	<u>SE</u>	<u>Wald</u>	<u>Step OR</u>	<u>Final OR</u>	<u>Final p</u>
1	Age	.44	.138	10.07	1.81	1.55	.002
	Gender ^a	-.42	.392	1.14	ns	ns	.286
	Race ^a	-1.10	.491	5.02	ns	3.01	.025
	Income	-.07	.094	0.60	0.82	ns	.439
2	Familial Problem Alc.	.87	.385	5.06	4.13	2.38	.025
	Familial Drug Use	.94	.392	5.74	4.12	2.56	.017
3	Physical Abuse/Asslt.	1.10	.459	5.78	4.68	3.01	.016
	Sexual Assault	1.00	.419	5.70	3.57	2.72	.017
4	Witnessed Violence	1.33	.584	5.21	3.99	3.79	.022
5	PTSD	.78	.430	3.30	ns	ns	.069

Note: Beta, SE, Wald, and p are given for the final model

^aOdds ratios greater than 1 for these variables indicate that being male or being Caucasian was associated with increased risk.

Table 10: Hierarchical Logistic Regression: Initial and Final Odds Ratio for Prediction of Past-year Delinquency for Male and Female Adolescents

<u>Prediction of Past-Year Delinquency: Male Adolescents</u>							
<u>Step</u>	<u>Variable</u>	<u>Beta</u>	<u>SE</u>	<u>Wald</u>	<u>Step OR</u>	<u>Final OR</u>	<u>Final p</u>
1	Age	.03	.05	.3	1.26	ns	.588
	Race*	.18	.18	1.0	0.59	ns	.314
	Income*	-.02	.04	.3	ns	ns	.561
2	Familial Problem Alc.	.60	.19	9.6	3.29	1.82	.002
	Familial Drug Use	.68	.23	8.9	3.16	1.97	.003
3	Physical Abuse/Asslt.	1.12	.16	47.2	4.84	3.06	.000
	Sexual Assault	.32	.32	1.0	1.84	ns	.316
4	Witnessed Violence	1.04	.18	33.4	3.20	2.82	.000
5	Lifetime PTSD	.40	.24	2.7	ns	1.50	.098
6	Problem Substance Use	1.49	.20	56.0	4.43	4.43	.000
<u>Prediction of Past-Year Delinquency: Female Adolescents</u>							
1	Age	-.08	.08	.9	1.22	0.40	.349
	Race*	.92	.27	11.6	0.47	ns	.001
	Income*	.05	.06	.5	ns	ns	.468
2	Familial Problem Alc.	.69	.27	6.5	3.60	1.99	.011
	Familial Drug Use	1.08	.28	15.1	4.33	2.93	.000
3	Physical Abuse/Asslt.	1.36	.28	23.0	6.61	3.88	.000
	Sexual Assault	.10	.29	.1	2.07	ns	.735
4	Witnessed Violence	1.34	.36	13.6	5.26	3.82	.000
5	Lifetime PTSD	.52	.29	3.3	1.73	1.68	.071
6	Problem Substance Use	1.80	.29	38.4	6.04	6.04	.000

*For these variables, odds ratios above 1 indicate that Caucasians are more likely than Non-Caucasians, and high income youth are more likely than low income youth to engage in delinquent behavior.

Table 11: Predictive Models of Past Year Index Offenses for White and African-American Males and Females

<i>Males</i>	White		African-American	
	n = 1,401		n = 277	
Variable	Step OR	Final OR	Step OR	Final OR
Low Income	1.92**	1.69	0.75	0.78
Family History Alcohol	3.22***	1.89*	2.74**	0.99
Family History Drug	4.69***	2.35**	1.19	0.84
Witnessed Violence	2.86***	2.15***	7.10***	5.96**
Sexual Assault	1.19	0.77	2.11	1.09
Physical Assault	3.68***	3.58***	2.71**	1.64
Physical Abuse	2.36**	1.96*	1.72	1.39
PTSD	1.37	1.58	3.16?*	3.26*
Significant Substance Use	4.03***	4.03***	6.65***	6.65***

<i>Females</i>	White		African-American	
	n = 1,345		n = 295	
Variable	Step OR	Final OR	Step OR	Final OR
Low Income	1.03	0.47	1.14	0.99
Family History Alcohol	5.92***	2.77**	4.84***	3.23*
Family History Drug	3.67***	1.96	3.29*	3.67*
Witnessed Violence	7.08***	5.18***	10.69**	9.16*
Sexual Assault	1.80	1.22	1.47	1.31
Physical Assault	3.34**	2.69*	3.35*	2.81*
Physical Abuse	0.62	0.62	2.24	2.12
PTSD	2.35*	1.93	1.26	1.31
Significant Substance Use	4.24***	4.24***	3.60*	3.60*

Table 12: Proportion of Male Youths Who Committed an Index Offense in the Past Year with Risk Factor Combination

Family History	Assault History	PTSD	Substance Abuse	% Delinquent	% Population	% Risk Comb.
-	-	-	-	20.0	61.2	4.5
+	-	-	-	3.6	5.7	8.7
-	+	-	-	21.0	13.5	21.3
-	-	+	-	2.2	1.4	20.7
-	-	-	+	5.4	3.5	21.2
+	+	-	-	12.9	5.2	34.3
+	-	+	-	1.3	0.7	27.0
+	-	-	+	3.3	0.6	78.4
-	+	+	-	1.6	1.5	14.0
-	+	-	+	8.2	2.5	44.5
-	-	+	+	0.0	0.1	0.0
+	+	+	-	3.2	0.9	49.5
+	+	-	+	8.6	1.7	71.2
+	-	+	+	0.0	0.1	0.0
-	+	+	+	2.2	0.5	58.2
+	+	+	+	6.5	1.0	89.9

Table 13: Proportion of Female Youths Who Committed an Index Offense in the Past Year with Risk Factor Combination

Family History	Assault History	PTSD	Substance Abuse	% Delinquent	% Population	% Risk Comb.
-	-	-	-	8.8	58.8	0.7
+	-	-	-	3.4	9.4	1.8
-	+	-	-	7.5	10.9	3.4
-	-	+	-	0.5	2.3	1.0
-	-	-	+	0.9	1.9	2.4
+	+	-	-	12.6	5.0	12.5
+	-	+	-	3.6	1.1	16.8
+	-	-	+	4.6	0.9	24.7
-	+	+	-	2.4	2.1	5.7
-	+	-	+	8.3	1.5	28.3
-	-	+	+	0.7	0.1	26.0
+	+	+	-	12.3	2.0	31.3
+	+	-	+	14.8	1.4	51.1
+	-	+	+	0.7	0.1	25.8
-	+	+	+	6.9	0.9	37.2
+	+	+	+	11.6	1.3	42.9

Table 14: Population Estimates of Victimization and PTSD

Variable	Cases in Sample	Prevalence in Sample	Population Estimate*
Sexual Assault	326	8.1%	1.8 million
Physical Assault	701	17.4%	3.9 million
Physically Abusive Punishment	376	9.4%	2.1 million
Witness Violence	1,586	39.4%	8.8 million
Lifetime PTSD	324	8.1%	1.8 million
Current PTSD	196	4.9%	1.1 million
Committed a Delinquent Offense Past Year	381	9.5%	2.1 million
Ever Committed a Delinquent Offense	496	12.3%	2.7 million
Current Alcohol Abuse/Dependence	157	3.9%	870 thousand
Current Marijuana Abuse/Dependence	147	3.7%	825 thousand
Current Any Substance Abuse/Dependence	277	6.9%	1.5 million
Lifetime Alcohol Abuse/Dependence	226	5.6%	1.3 million
Lifetime Marijuana Abuse/Dependence	179	4.5%	1 million
Lifetime Drug Abuse/Dependence	46	1.2%	268 thousand
Lifetime Any Substance Abuse/Dependence	366	9.1%	2.0 million

*Based on Bureau of Census 1995 estimates that U.S. population of adolescents is 22.3 million. Rounded to nearest 100,000.

Figure 1.
Age at Time of Sexual Assault (n=462 cases)




Figure 2. Relationship Between Victim and Perpetrator in Sexual Assault Cases (n=462 cases)




Figure 2 (continued).
Relationship Between Victim and Perpetrator in Sexual
Assault Cases (n=462 cases)





Figure 3.
Location of Sexual Assaults (n=462 cases)



**Figure 4 .
 Life Threat and Degree of Physical Injury
 Sustained During Sexual Assault (n=462 cases)**




Figure 5.
Reporting of Sexual Assault to Authorities
(n=462 cases)




Figure 6.
Victims Age at Time of Physical Assault
(n=1,054 cases)




Figure 7.
Relationship Between Physical Assault Victims and their Perpetrators (n=1,054 cases)




Figure 8.
Location of Physical Assault (n=1,054 cases)




Figure 9.
Life Threat and Degree of Physical Injury
Sustained During Physical Assault (n=1,054 cases)




Figure 10.

Reporting of Physical Assault to Authorities (n=1,054 cases)




Figure 11.

Lifetime Prevalence of Witnessing Violence





Figure 12.
Lifetime Prevalence of Sexual Assault
by Age Cohort



RR

Figure 13. Lifetime Prevalence of Physical Assault by Age Cohort




Figure 14.

Victims Age at Time of Physically Abusive Punishment by Age Cohort




Figure 15.
Age at Time Witnessed Violence by Age Cohort




Figure 16.
Prevalence Rates of Lifetime and Current PTSD by Age Cohort




Figure 17.
Lifetime and Past Year Heavy Alcohol Use
by Age Cohort




Figure 18.
Lifetime and Past Year Use of Illicit Drugs by Age Cohort




Figure 19.

Past Year Rates of Alcohol, Marijuana, and Hard Drug Abuse/Dependence




Figure 20.

Lifetime and Past Year Delinquent Offense by Age Cohorts




Figure 21.

Percentage of Male and Female Adolescents with Lifetime PTSD by Number of Sexual Assaults Experienced




Figure 22.

Percentage of Male and Female Adolescents with Lifetime PTSD by Number of Physical Assaults Experienced (N=4,023)




Figure 23.

Percentage of Male and Female Adolescents with Lifetime PTSD by Number of Incidents of Violence Witnessed (N=4,023)





Figure 24.

Rates of Past Year Any Substance Abuse/Dependence by Victimization Risk Factors and Gender (N=4,023)



(X)

Figure 25.
Past Year Delinquency by Victimization Risk Factors
and Gender (N=4,023)




Figure 26.

Percentage of Male and Female Adolescents with PTSD by Family Members with Alcohol and Drug Problems (N=4,023)





Figure 27.

Number of Violent Incidents Experienced or Witnessed by Male and Female Adolescents (N=4,023)

Male Adolescents



Female Adolescents




Figure 28.

Past Year Alcohol Abuse/Dependence (N=4,023)




Figure 29.
Past Year Marijuana Abuse/Dependence (N=4,023)

